



## 1 Decay Scheme

Te-123m decays via two successive gamma transitions.

A gamma transition with a small probability and an energy of 247 keV has been observed.

*Te-123m se désintègre via deux transitions gamma en cascade. Une transition de 247 keV et de faible probabilité a été observée.*

## 2 Nuclear Data

$$\begin{aligned}
 T_{1/2}(^{123}\text{Te}^m) &: 119,3 \quad (1) \quad \text{d} \\
 T_{1/2}(^{123}\text{Te}) &: 12 \quad 10^{12} \text{ a}
 \end{aligned}$$

### 2.1 Gamma Transitions and Internal Conversion Coefficients

	Energy keV	$P_{\gamma+ce}$ $\times 100$	Multipolarity	$\alpha_K$	$\alpha_L$	$\alpha_M$	$\alpha_T$
$\gamma_{2,1}(\text{Te})$	88,46 (7)	100,0 (1)	M4	463 (14)	493 (15)	118,0 (35)	1099 (33)
$\gamma_{1,0}(\text{Te})$	158,99 (5)	100,0 (1)	M1+1,22%E2	0,1648 (16)	0,02160 (22)	0,00433 (13)	0,1918 (19)
$\gamma_{2,0}(\text{Te})$	247,4 (2)	0,0030 (3)	E5	3,0 (1)	3,75 (21)	0,84 (3)	7,75 (30)

## 3 Atomic Data

### 3.1 Te

$$\begin{aligned}
 \omega_K &: 0,875 \quad (4) \\
 \bar{\omega}_L &: 0,0862 \quad (35) \\
 n_{KL} &: 0,917 \quad (4)
 \end{aligned}$$

**3.1.1 X Radiations**

	Energy keV	Relative probability
$X_K$	$K\alpha_2$	27,202
	$K\alpha_1$	27,4726
	$K\beta_3$	30,9446
	$K\beta_1$	30,996
	$K\beta_5''$	31,236
	$K\beta_5'$	31,241
	$K\beta_2$	31,7008
	$K\beta_4$	31,774
	$KO_{2,3}$	31,812
$X_L$	$L\ell$	3,336
	$L\alpha$	3,76 – 3,77
	$L\eta$	3,606
	$L\beta$	4,02 – 4,37
	$L\gamma$	4,44 – 4,82

**3.1.2 Auger Electrons**

	Energy keV	Relative probability
Auger K		
KLL	21,804 – 22,989	100
KLX	25,814 – 27,470	45,3
KXY	29,80 – 31,81	5,13
Auger L		
	2,3 – 4,8	

## 4 Electron Emissions

		Energy keV	Electrons per 100 disint.
e <sub>AL</sub>	(Te)	2,3 - 4,8	89,7 (7)
e <sub>AK</sub>	(Te)		7,0 (4)
	KLL	21,804 - 22,989	}
	KLX	25,814 - 27,470	}
	KXY	29,80 - 31,81	}
ec <sub>2,1</sub> K	(Te)	56,65 (7)	42,1 (18)
ec <sub>2,1</sub> L	(Te)	83,52 - 84,12	44,8 (19)
ec <sub>2,1</sub> M	(Te)	87,45 - 87,89	10,73 (45)
ec <sub>2,1</sub> N	(Te)	88,29 - 88,42	2,07 (9)
ec <sub>1,0</sub> K	(Te)	127,18 (5)	13,84 (14)
ec <sub>1,0</sub> L	(Te)	154,05 - 154,69	1,81 (2)
ec <sub>1,0</sub> M	(Te)	157,98 - 158,42	0,364 (11)
ec <sub>1,0</sub> N	(Te)	158,82 - 158,95	0,0769 (23)

## 5 Photon Emissions

### 5.1 X-Ray Emissions

		Energy keV	Photons per 100 disint.	
XL	(Te)	3,336 — 4,82	8,25 (21)	
XK $\alpha_2$	(Te)	27,202	13,9 (5)	} K $\alpha$
XK $\alpha_1$	(Te)	27,4726	26,0 (9)	}
XK $\beta_3$	(Te)	30,9446	}	
XK $\beta_1$	(Te)	30,996	}	K' $\beta_1$
XK $\beta_5''$	(Te)	31,236	}	
XK $\beta_5'$	(Te)	31,241	}	
XK $\beta_2$	(Te)	31,7008	}	
XK $\beta_4$	(Te)	31,774	}	K' $\beta_2$
XKO <sub>2,3</sub>	(Te)	31,812	}	

## 5.2 Gamma Emissions

	Energy keV	Photons per 100 disint.
$\gamma_{2,1}(\text{Te})$	88,46 (7)	0,0909 (27)
$\gamma_{1,0}(\text{Te})$	158,97 (5)	83,99 (8)
$\gamma_{2,0}(\text{Te})$	247,4 (2)	0,000344 (34)

## 6 Main Production Modes

- $\left\{ \begin{array}{l} \text{Te} - 122(\text{n}, \gamma) \text{Te} - 123\text{m} \quad \sigma : 3,4 (5) \text{ barns} \\ \text{Possible impurities : Te} - 121, \text{Te} - 125\text{m}, \text{Te} - 129\text{m} \end{array} \right.$
- $\left\{ \begin{array}{l} \text{Sb} - 123(\text{p}, \text{n}) \text{Te} - 123\text{m} \\ \text{Possible impurities : Te} - 121, \text{Te} - 121\text{m} \end{array} \right.$
- $\left\{ \begin{array}{l} \text{Sb} - 123(\text{d}, 2\text{n}) \text{Te} - 123\text{m} \\ \text{Possible impurities : Te} - 121, \text{Te} - 121\text{m} \end{array} \right.$

## 7 References

- L.W.FAGG, E.A. WOLIKI, R.O. BONDELID, K.L. DUNNING, S. SNYDER. Phys. Rev. 100 (1955) 1299 (Mixing ratio)
- N.GOLDBERG, S.FRANKEL. Phys. Rev. 100 (1955) 1350 (Mixing ratio)
- Y.Y.CHU, M.L.PERLMAN. Phys. Rev. 135,2B (1964) 319 (ICC)
- Y.Y.CHU, O.C.KRISTNER, A.C.LI, S.MONARO, M.L.PERLMAN. Phys. Rev. 133,6B (1964) 1361 (ICC, Gamma-ray intensities)
- D.G.ALKHAZOV, V.D. VASILEV, G.M. GUSINSKII, I.K. LEMBERG, V.A. NABICHVRISHVILI. Bull. Acad. Sci. USSR Phys. Serv. 28 (1964) 1575 (Mixing ratio)
- E.N.HATCH, G.W.EAKINS, G.C.NELSON, R.E.MC ADAMS. Proc. Intern. Conf. Intern. Conversion Processes, Nashville, Tenn. (1965), Ed. J.H.Hamilton (1966) 183 (Gamma-ray emission probabilities, ICC)
- R.K.GUPTA, M.M. BAJAJ, N.K. SAHA. Nucl. Phys. 80 (1966) 471 (Mixing ratio)
- S.TÖRNKVIST, S.STROM, L.HASSELGREN. Nucl. Phys. A130 (1969) 604 (Mixing ratio)
- R.A.KALINAUSKAS, K.V.MAKARYUNAS, K.MAKATYUNENE, R.I.DAVIDONIS. Bull. Acad. Sci. USSR Phys. Serv. 32 (1969) 187 (T ICC)
- J.F.EMERY, S.A.REYNOLDS, E.I.WAYTT. report ORNL-4466 (1970) 75 (Half-life)
- S.RAMAN. Nucl. Instrum. Methods 103 (1972) 407 (Gamma-ray emission intensities)
- S.RAMAN, R.L.AUBLE, W.T.MILNER, T.A.WALKIEWICZ, R.GUNNINK, B.MARTINE. Report ORNL-4937 (1973) 144 (ICC)

- S.RAMAN, R.L.AUBLE, W.T.MILNER. Phys. Lett. 47B (1973) 19  
(Conv. Elec. emission probabilities, Gamma-ray emission probabilities, Multipolarity)
- K.S.KRANE. At. Data. Nucl. Data Tables 19 (1977) 19  
(Mixing ratio)
- W.BAMBYNEK. X-84 Proc. X-Ray and Inner-Shell Processes in Atoms, Molecules and Solids, A. Meisel Ed., Leipzig Aug. 20-23 (1984)  
(K fluorescence yield)
- U.SCHÖTZIG, H.SCHRADER, K.DEBERTIN. Proc. Int. Conf. Nuclear Data Sci. Techn., Jülich, 13-17 May (1991)  
(Gamma-ray emission probabilities)
- B.M.COURSEY, D.B.GOLAS, D.H.GRAY, D.D.HOPPES, F.J.SCHIMA. Nucl. Instrum. Methods A312 (1992) 121  
(Gamma-ray emission probability, Half-life)
- H.JANSSEN, E.SCHÖNFELD, R.KLEIN. Int. J. Appl. Radiat. Isotop. 43,11 (1992) 1309  
(T ICC (1,0), Gamma-ray emission probability)
- E.SCHÖNFELD, H.JANSSEN. Nucl. Instrum. Methods A369 (1996) 527  
(Atomic Data)
- I.M.BAND, M.B.TRZHASKOVSKAYA. At. Data. Nucl. Data Tables 88,1 (2002)  
(Theoretical ICC)
- V.M.GOROZHANKIN, N.COURSOL, E.A.YAKUSHEV, Ts.VYLOV, C.BRIANÇON. Appl. Rad. Isotopes 56 (2002) 189  
(M4 transition)

$\gamma$  Emission probabilities  
per 100 disintegrations

