


Table de RADIONUCLÉIDES

Table of RADIONUCLIDES

Tabelle der RADIONUKLIDE



INTRODUCTION

Note technique LNHB 2011/53

Marie-Martine BÉ, Nelcy COURSQL, Bernard DUCHEMIN, Frédérick LAGOUTINE, Jean LEGRAND
(CEA, LIST, LNE-LNHB)

Klaus DEBERTIN, Eckart SCHÖNFELD, Physikalisch-Technische Bundesanstalt (PTB)

« Table de RADIONUCLÉIDES – Introduction »

Sommaire

Ce fascicule est une introduction à la Table de Radionucléides publiée sur papier et également à sa version informatisée NUCLÉIDE disponible sur CD-Rom. Il inclut une brève description des processus physiques de la radioactivité, l'énumération des règles d'évaluation conduisant aux valeurs recommandées, ainsi qu'un récapitulatif des symboles et des termes utilisés dans les différentes publications.

“Table of RADIONUCLIDES – Introduction”

Summary

This volume is an introduction to the Table of Radionuclides published on paper and also to its computerized form NUCLÉIDE available on CD-Rom. It includes a brief description of the radioactivity physical processes, the enumeration of the evaluation rules leading to the recommended values, and a summary of the symbols and terms used in all the publications.

„Tabelle der RADIONUKLIDE – Einführung“

Zusammenfassung

Dieser Text ist eine Einführung zur Tabelle der Radionuklide, in gedruckter Form ebenso wie zur auf CD-ROM erhältlichen elektronischen Ausgabe NUCLÉIDE. Er umfasst eine kurze Beschreibung der physikalischen Prozesse der Radioaktivität, eine Auflistung der Auswertungsregeln, die zu den empfohlenen Werten führen, und eine Zusammenfassung der in den diversen Veröffentlichungen verwendeten Symbole und Begriffe.

Table de RADIONUCLÉIDES
Table of RADIONUCLIDES
Tabelle der RADIONUKLIDE

INTRODUCTION

Edition 2011 : Marie-Martine BÉ (CEA/LIST – LNE/LNHB)

Edition 1999 : Marie-Martine BÉ, Bernard DUCHEMIN (BNM-CEA/LPRI)
Klaus DEBERTIN, Eckart SCHÖNFELD, Physikalisch-Technische Bundesanstalt (PTB)

Edition 1987 : Frédérick LAGOUTINE, Nelcy COURSOLO, Jean LEGRAND

Contents

1	PREFACE	3
2	SYMBOLS AND NOTATIONS	5
2.1	Units	5
2.2	Particles and quanta	5
2.3	Energies	6
2.4	Transition probabilities, emission intensities and conversion coefficients	7
2.5	Other physical quantities and abbreviations	8
3	DESCRIPTION OF RADIOACTIVE PROCESSES	10
3.1	Brief reminder of some definitions	10
3.2	Disintegration modes	13
3.2.1	α decay	13
3.2.2	β decay	15
3.2.3	Electron capture decay	19
3.2.4	ϵ/β^+ ratio	22
3.3	γ transitions	22
3.3.1	γ photon emission	22
3.3.2	Internal conversion	23
3.3.3	Internal pair creation	24
3.3.4	Multipolarity	24
3.3.5	Mean life times of excited levels	25
3.4	Secondary phenomena accompanying nuclear transformations	26
3.4.1	Internal bremsstrahlung accompanying β disintegration	26
3.4.2	Internal bremsstrahlung accompanying electron capture	26
3.4.3	Internal ionization and excitation accompanying β decay	27
3.4.4	Internal ionization and excitation accompanying electron capture	27
3.4.5	Other processes	28
3.5	Electron rearrangement	28
3.5.1	K shell	28
3.5.2	L shell	30
4	INTERACTION OF RADIATIONS WITH MATTER	35
4.1	External bremsstrahlung	35
4.2	Annihilation-in-Flight of Positrons	35
5	DOSIMETRIC DATA	36

6	RULES FOR EVALUATION AND COMPILATIONS	38
6.1	Rules for evaluation	38
6.1.1	Evaluation of uncertainties	39
6.1.2	Determination of the best value and of the associated uncertainty	39
6.1.3	Balanced decay schemes	42
6.2	Compilations	43
6.2.1	β and electron capture transitions	43
6.2.2	γ transitions	43
6.2.3	Level spins and parities	44
6.2.4	Constants of the atomic shell	44
6.2.5	m_0c^2 energy	44
7	CONTENTS OF THE TABLES	45
7.1	Radioactive series	45
7.2	Decay scheme	45
7.3	Tables	47
7.3.1	Notation	47
7.3.2	Nuclear data	47
7.3.3	Atomic data	48
7.3.4	Intensities of radiations	48
7.4	Main production modes	49
7.5	References used	49
7.6	Spectra	49
7.7	Publications	49
	REFERENCES	50

Chapter 1

PREFACE

The aim of this Table is to provide carefully evaluated data for radionuclides frequently used in science and technology. The evaluation is based on presently available experimental data and, in part, on theoretical consideration and the finally evaluated (and recommended) data are presented in a clear and easily readable form. This publication is a continuation of the earlier “Table de Radionucléides” (LMRI, France, 1974-1980), by F. Lagoutine, N. Coursol and J. Legrand (LMRI, France, 1982-1986).

In order to update the data of the nuclides already present and to add new evaluations, the Laboratoire National Henri Becquerel (LNHB, France) and the Physikalisch-Technische Bundesanstalt (PTB, Germany) established a cooperative agreement; they were then joined by the Idaho National Engineering & Environmental Laboratory (INEEL, USA), the Lawrence Berkeley National Laboratory (LBNL, USA) and the Khlopin Radium Institute (KRI, Russia). This international collaboration is based on an informal agreement, the initial work of this group was to discuss and to agree on a methodology to be used in these evaluations. This is described in the chapter “Rules for Evaluation and Compilation”.

In this new version, there are small changes in the presentation following international recommendations (e.g. use of the standard deviation instead of the uncertainty at the 99,7 % confidence level which was used before 1986) but the main objectives and contents are the same.

Over the past years, several volumes of data and comments on how the recommended data have been obtained were published as CEA Reports [99 TR2] or as BIPM Monographie [04TRM1, 06TRM2, 08TRM3, 10TRM4, 11TRM5].

New available evaluations of radionuclides are updated on:

<http://www.nucleide.org/NucData.htm>

Moreover, LNHB developed a software NUCLÉIDE [CD NUC] with the objectives of making it easier to update and add data and, obviously, to offer easy access to the nuclear and atomic decay data to the user by “button click” facilities.

Each evaluated radionuclide is presented in a separate section, which can be practical for users and facilitates updating if new experimental work makes this necessary.

Each section includes:

- a decay scheme with the minimum information required for understanding the involved processes and levels;
- tables listing the nuclear data and atomic data directly relevant to the main emitted radiations;

- tables of intensities for emission of alpha and electron particles and photon radiations which, in addition to the half-life, are the most useful data from the practical standpoint;
- the most important production method(s) and potential radioactive impurities. When one of these production modes involves activation by thermal neutrons, the cross section is given;
- the references employed with indications of parameters used from each. All references known by the author concerning a radionuclide have been examined, but only those employed in the evaluation are listed;
- Finally the evaluation date and the author of the evaluation, who can provide any additional information, are given.

Système International (SI) units [81 SI], ISO 31 recommendations [92 ISO] and Guide for the Expression of Uncertainty in Measurement [93 GUM] are used throughout in this text.

Chapter 2

SYMBOLS AND NOTATIONS

2.1 Units

s	second
min	minute
h	hour
d	day
a	year (1 a = 365,242 198 78 d or 31 556 925,26 s)
m	meter
J	joule
eV	electronvolt (1 eV = 1,602 176 462 (63) $\times 10^{-19}$ J)
Bq	becquerel (1 Bq = 1 disintegration.s ⁻¹)
Δ	dose constant (J.Bq ⁻¹ .s ⁻¹)

2.2 Particles and quanta

e	electron
p	proton
n	neutron
d	deuteron
α	alpha particle
β^+	positron from β^+ decay
β^-	electron from β^- decay
γ	gamma quantum, photon emitted when a nucleus turns into a lower energy state
ν	neutrino ($\bar{\nu}$ antineutrino)
ce	internal conversion electron
ce _K	internal conversion electron, ejected from the K shell
ce _L	internal conversion electron, ejected from the L shell
e _A	Auger electron
e _{AK}	K-Auger electron
e _{AL}	L-Auger electron
KLL	KLL-Auger electron

KLX	KLX-Auger electron (X=M,N)
KXY	KXY-Auger electron (X=M,N; Y=M,N)
X	x-ray quantum, photon emitted during the rearrangement of the atomic shells
X _K	K X-ray quantum
X _L	L X-ray quantum
e [±]	internal electron-positron pair
γ [±]	gamma photon due to positron annihilation

2.3 Energies

E	energy, generally
E_{α}	alpha particle energy
$E_{\beta^{+max}}$	β^{+} particle maximum energy
$E_{\beta^{-max}}$	β^{-} particle maximum energy
$\bar{E}_{\beta^{+}}$	β^{+} particle mean energy
$\bar{E}_{\beta^{-}}$	β^{-} particle mean energy
E_{γ}	gamma quanta energy
E_{ph}	energy of photons which are not gamma quanta with regard to their origin
E_r	recoil energy
E_i	energy of a nucleus excited level
E_K	binding energy of an electron in the K shell
E_L	binding energy of an electron in the L shell
E_n	neutron energy
E_{ν}	neutrino energy
E_e	monoenergetic electron energy
E_{AK}	K-Auger electron energy
E_{AL}	L-Auger electron energy
E_{XK}	K X-ray quanta energy
E_{XL}	L X-ray quanta energy
E_{ceK}	K conversion electron energy
E_{ceL}	L conversion electron energy
Q_{α}	total available energy of alpha disintegration
$Q_{\beta^{+}}$	total available energy of β^{+} disintegration
$Q_{\beta^{-}}$	total available energy of β^{-} disintegration
Q_{ϵ}	total available energy of electron capture transformation
q_K	energies of neutrinos after electron capture in the K shell
q_L	energies of neutrinos after electron capture in the L shell
q_M	energies of neutrinos after electron capture in the M shell
W	total energy of a β particle including its rest energy

2.4 Transition probabilities, emission intensities and conversion coefficients

P_α	transition probability for an alpha transition
P_β	transition probability for a beta transition
P_e	monoenergetic electron emission probability
P_ϵ	transition probability for an electron capture transition
P_K	K-shell capture probability for an Electron Capture (EC) transition
P_L	L-shell capture probability for an EC transition
P_M	M-shell capture probability for an EC transition
	$(P_K + P_L + P_M + \dots = 1)$
P_g	Total transition probability for a gamma transition including conversion electrons
	$P_g = I_\gamma + I_{ce}$
I_γ	gamma emission intensity
I_{ce}	conversion electron emission intensity
I_{ceK}	K conversion electron emission intensity
I_{ceL}	L conversion electron emission intensity
I_{ceM}	M conversion electron emission intensity
I_X	X ray emission intensity
I_{XK}	K X-ray emission intensity
I_{XL}	L X-ray emission intensity
I_{XM}	M X-ray emission intensity
I_A	Auger electron emission intensity
I_{AK}	K-Auger electron emission intensity
I_{AL}	L-Auger electron emission intensity
I_{AM}	M-Auger electron emission intensity
I_{AKLL}	KLL-Auger electron emission intensity
λ_n	probability for a given β transition
λ_0	probability for an allowed β transition
α_K	K-shell internal conversion coefficient
α_L	L-shell internal conversion coefficient
α_M	M-shell internal conversion coefficient
α_t	total internal conversion coefficient ($\alpha_K + \alpha_L + \alpha_M + \dots = \alpha_t$)
α_π	internal pair creation coefficient

2.5 Other physical quantities and abbreviations

a_K	K-Auger yield, $a_K = 1 - \omega_K$
\bar{a}_L	mean L-Auger yield, $\bar{a}_L = 1 - \bar{\omega}_L$
A	(1) activity (2) mass number in A_ZX_N , where X is an element
B_K, B_{L_1}	exchange factor in the K shell, L_1 subshell,...
C_n	shape factor of a β transition
C_0	shape factor of an allowed β transition
E0, E1, E2, EL	electric monopole, dipole, quadrupole, 2L-pole
f	(1) component of radial functions of the electron (2) alpha particle collision frequency
f_0	Fermi integral for an allowed transition
f_1, f_2, \dots	Fermi integrals for unique 1 st forbidden, 2 nd forbidden,... transitions
ft	comparative half-life
f_{12}, f_{13}, f_{23}	probabilities of Coster-Kronig transitions
F	hindrance factor
$F(t)$	correction factor for radioactive decay
g	(1) coupling constant of weak interactions (2) components of radial functions of the electron
J	quantum number of total angular momentum
K, L, M ...	electron shells
K/L	ratio $I_{ceK}/I_{ceL} = \alpha_K/\alpha_L$
K/LM	ratio $I_{ceK}/(I_{ceL} + I_{ceM}) = \alpha_K/(\alpha_L + \alpha_M)$
K/LMN	ratio $I_{ceK}/(I_{ceL} + I_{ceM} + I_{ceN}) = \alpha_K/(\alpha_L + \alpha_M + \alpha_N)$
KLX/KXY	ratio I_{AKLX}/I_{AKXY}
L	orbital angular momentum quantum number
L_1, L_2, L_3	L electron subshells
m_0	electron rest mass
max	maximum
min	minimum
av	average
M	atomic mass
M_N	mass of the nucleus
M1, M2, ML	magnetic dipole, quadrupole, 2L-pole
n_K, n_L	mean number of vacancies in the K shell and L shell created during the disintegration of one nucleus of a given radionuclide
n_{KL}	total number of L vacancies created by $K \rightarrow L$ transfer, i.e. mean number of vacancies in the L shell produced by one vacancy in the K shell
N	(1) number of neutrons in A_ZX_N where X is an element (2) number of radioactive atoms
P	penetrability for the potential barrier
R	radius of the nucleus

r_0	radius of the nucleus in units of \hbar/m_0c
S	intrinsic spin
t	time
$T_{1/2}$	radioactive half-life
v_0	velocity of the α particle
V_{L_i}	Coster-Kronig vacancies
Z	atomic number in A_ZX_N where X is an element
δ	ratio $\delta^2 = E_{(L+1)}/M_{(L)}$
	mixing of different multipolarities
ϵ	transition by electron capture (EC)
λ	decay constant
π	parity
σ	total neutron capture cross section
σ^0	2200 m.s ⁻¹ neutron capture cross section
ω_K	K-shell fluorescence yield
$\omega_{L1}, \omega_{L2}, \omega_{L3}$	fluorescence yield in the L ₁ , L ₂ , L ₃ subshells
$\bar{\omega}_L$	mean L-shell fluorescence yield
$\bar{\omega}_M$	mean M-shell fluorescence yield

Chapter 3

DESCRIPTION OF RADIOACTIVE PROCESSES

3.1 Brief reminder of some definitions

Nuclide: A nuclide is a species of atoms characterized by an atomic number Z (proton number) and a mass number A (nucleon number).

Radionuclide: A radionuclide is a radioactive nuclide.

Disintegration: A disintegration of an atomic nucleus is the spontaneous transformation of the nucleus into another one giving rise to a change in the atomic number, or into two or more nuclei (fission), or, in the case of an isomer, a transition to a lower energy state of the same nucleus. A disintegration leaves the daughter nucleus in the ground state or in an excited state ; a de-excitation occurs through one or more γ transitions. While in α decay there is also a change in the mass number, in β^- or β^+ decay and electron capture decay the mass number remains unchanged.

Transition: A transition is a process in which an atomic nucleus undergoes a transformation by disintegration or de-excitation.

Isomer: An isomer is a nuclide in an excited state having a non-negligible half-life, liable to de-excitation by a γ transition (isomeric transition, e.g. $^{103\text{m}}\text{Rh}$), or to any other transition, such as β decay or electron capture (e.g. $^{110\text{m}}\text{Ag}$, $^{115\text{m}}\text{In}$).

Transition probability: The transition probability assigned to a specific transition is the probability that this transition occurs when one nucleus of this radionuclide disintegrates.

The transition probability includes all processes which lead from the initial level to the final level, where the latter may belong to the daughter nucleus or to the same nucleus.

With respect to different transition types, the transition probability is the probability of the transformation of a nuclide in a given energy state *via* the emission of particles (α , β^- , β^+) or the capture of electrons from the atomic shell (denoted by ϵ or EC) into a specific state of their daughter nuclei (excited state or ground state) or *via* the emission of gamma photons and conversion electrons into a lower energy state of nuclei of the same nuclide.

According to its above definition the probability of a transition has a value between 0 and 1. A radionuclide may decay by one type of transitions only or it may have branches of different types (for example EC and β^+). In each case the sum of the transition probabilities of all branches to the daughter nuclide(s) is exactly 1.

Emission intensity: The emission intensity assigned to the particles or quanta (specified by type and energy) emitted by a radionuclide is the mean number of particles or quanta emitted per disintegration of one nucleus. The emission intensities are generally related to 100 nuclear transformations of the studied nuclide.

For X-rays, Auger electrons and γ emissions due to positron annihilation, the emission intensity may exceed 1 due to the following reasons:

- vacancies in the K-shell and other shells can be produced by EC transitions as well as by inner conversion of γ -transitions;
- the number of vacancies in the L(M) shell, created by one vacancy in the K(L) shell can be larger than 1;
- two photons are emitted per one emitted positron.

For these reasons, in the tables, the notation “number of electrons (or photons or alpha) per disintegration” are used throughout.

The following relationships state between transition probabilities and emission intensities:

- an α transition probability is equal to the mean number of α particles emitted per disintegration (or α emission intensity);
- a β transition probability is equal to the mean number of β particles emitted per disintegration (or β emission intensity);
- an EC transition probability is composed by the contributions of capture processes in the K, L, M, ... shells. It is related to the emission intensities for X rays and Auger electrons *via* fluorescence yields and other atomic shell data;
- a γ transition probability is the sum of the mean number of photons (or photon emission intensity) and conversion electrons (or conversion electron emission intensity) from different shells (K, L, M, ...) emitted per disintegration.

Energy of disintegration: The energy of disintegration is the available energy from the ground state (plus energy of the excited level in the case of isomers) of the parent to the final state of the daughter (in most cases the ground state level). It corresponds to the difference in atomic masses between these levels.

Activity: The activity, A , of an amount of radioactive nuclide in a particular energy state at a given time is the quotient of dN by dt , where dN is the expected value of the number of spontaneous nuclear transitions from that energy state in the time interval dt :

$$A = \frac{dN}{dt} \quad (3.1)$$

(Definition according to the International Commission on Radiation Units and Measurements (ICRU) [11ICRU]. The “particular energy state” is the ground state of the nuclide unless otherwise specified, e.g. in the case of isomers.

The special name for the unit of activity is **becquerel (Bq)**, 1 Bq = one disintegration per second.

The old unit of activity, the *curie* (Ci), is still used in some medical circles; its value is $1 \text{ Ci} = 37 \times 10^9 \text{ Bq}$.

Radioactive decay: The radioactive decay is the decrease, with time, of the activity of a radioactive substance by spontaneous nuclear transformations. It is expressed by the equations:

$$\frac{dN}{dt} = -\lambda N \quad (3.2)$$

$$N = N_0 e^{-\lambda t} \quad (3.3)$$

$$A = A_0 e^{-\lambda t} \quad (3.4)$$

where N is the number of atoms of the corresponding radionuclide consisting of nuclei being in the same energy state at time t , N_0 is the number of such atoms at time $t = 0$ and λ is the decay constant. Analogously, A and A_0 are the activity of the radioactive source or substance at time t and at time $t = 0$. When several nuclear transformation modes are in competition with partial decay constants λ_i (e.g. $^{110\text{m}}\text{Ag}$, ^{58}Co , ^{186}Re , etc.), the decay constant λ is written:

$$\lambda = \sum \lambda_i \quad (3.5)$$

The half-life $T_{1/2}$ is the time required for the initial number of atomic nuclei, in a given state, to decrease by a factor of two.

$$T_{1/2} = \frac{\ln 2}{\lambda} \quad (3.6)$$

The mean life or expected life-time $t = 1/\lambda$ is frequently employed.

Remark: in practice, we advise against using the expression “partial half-life” defined by $(T_{1/2})_i = \ln 2/\lambda_i$, which has no practical meaning.

One such example is ^{252}Cf , with a half-life of 2,6 years, which can disintegrate by α emission or by spontaneous fission and, which therefore has two partial decay constants λ_α and λ_F . Then:

$$\lambda = \lambda_\alpha + \lambda_F \quad (3.7)$$

The activity of a source of ^{252}Cf is given by equations (3.2) and (3.4). The α emission rate is $\lambda_\alpha N$ and that of spontaneous fission is $\lambda_F N$, each of these rates decreasing with $e^{-\lambda t}$ and not with $e^{-\lambda_\alpha t}$ or $e^{-\lambda_F t}$. Then:

$$T_{1/2} = \frac{\ln 2}{\lambda_\alpha + \lambda_F} \quad (3.8)$$

$$T_{1/2} = \frac{(T_{1/2})_\alpha \times (T_{1/2})_F}{(T_{1/2})_\alpha + (T_{1/2})_F} \quad (3.9)$$

The decay constants and half-lives are, in principle, independent of extranuclear conditions. In fact, radioactivity is not a purely nuclear phenomenon [$^{73}\text{Cr B}$]. Changes occurring in the environment, more precisely in the electrons of the external shells, can

cause decay constant variations [72Em G]. The different types of nuclear transformation able to give rise to observable changes are those in which the orbital electrons contribute directly to the nuclear transformation.

In the case of ${}^7\text{Be}$, which disintegrates by electron capture, differences, $\Delta\lambda/\lambda$, greater than 10^{-3} have been observed between various chemical compounds. In the case of ${}^{99\text{m}}\text{Tc}$, partly de-excited by a strongly converted γ transition of 2,17 keV, differences of the order of $0,3 \times 10^{-3}$ have also been observed.

3.2 Disintegration modes


Only α , β^- , β^+ and ϵ (electron capture) decays are examined here. The other existing modes of disintegration, such as spontaneous fission, proton emission and delayed neutron emission, will be examined later, if necessary, in an appendix.

Nuclei of a given radionuclide can disintegrate *via* a single transition mode, for example by β^- disintegration only or *via* several transition modes which are in competition with each other, such as β^+ and electron capture or β^- and α .

Isomers can de-excite or undergo radioactive disintegration (e.g. ${}^{110\text{m}}\text{Ag}$, ${}^{115\text{m}}\text{In}$).

3.2.1 α decay

A nucleus with atomic number Z and mass number A disintegrates by emitting an α particle, and giving a nucleus with atomic number $Z - 2$ and mass number $A - 4$;



The created nucleus may be in an excited energy state E_i .

Hence the α disintegration energy is:

$$Q_\alpha = E_{\alpha_i} + E_i + E_{r_i} \quad (3.10)$$

where E_{α_i} is the energy of the emitted α particle, constant for a given transition, and E_{r_i} the recoil energy:

$$E_{r_i} = \frac{M_\alpha}{M_N + M_\alpha} \left[Q_\alpha - E_i \right] \quad (3.11)$$

with M_N : mass of the recoiling daughter nucleus and M_α : mass of the α -particle.

Using (3.10) and (3.11):

$$E_{r_i} = \frac{M_\alpha}{M_N} E_{\alpha_i} \quad (3.12)$$

Probability of α decay

This is determined by means of the theory of the one-body model. When an α particle moves within a spherical nucleus and is held there by the Coulomb potential barrier, a certain probability exists that the α particle will escape from the nucleus, although the potential energy represented by the height of the barrier is greater than its kinetic energy (tunneling).

The decay constant for a given α transition is expressed by:

$$\lambda = f P = \frac{V_0}{R} P \quad (3.13)$$

where:

- f : collision frequency of the α particle within the potential well,
- V_0 : velocity of the α particle within this potential well,
- R : radius of the nucleus,
- P : penetrability of the potential barrier.

P is a function of the atomic number, the mass number, the radius of the nucleus, and the characteristics of the potential barrier.

The α transitions of even-even nuclei from ground state to ground state have the highest decay constants. They are adequately described by theoretical expressions based on (3.12), or by the following equation which is derived from (3.12), for a given atomic number:

$$\lg (T_{1/2})_i = AQ^{-1/2} + B \quad (3.14)$$

where:

- $(T_{1/2})_i$: partial half-life of the transition,
- Q : total energy available for this transition,
- A and B : constants calculated on the basis of experimental values of Q and transition probabilities from ground state to ground state.

The theory of the one-body model completely ignores the probability of preformation of the α particle, as well as the effect of the angular momentum carried off by the α particle on the penetrability of the potential barrier. This means that the experimental values of partial half-lives other than those of even-even nucleus transitions from ground state to ground state, are generally greater than calculated values. They are hindered at various degrees [64Hy E, 68Ra J].

Hindrance factor

The hindrance factor F for a given α transition is defined by the ratio:

$$F = \frac{(T_{1/2})_{i \text{ exp}}}{(T_{1/2})_{i \text{ th}}} \quad (3.15)$$

where:

$(T_{1/2})_{i \text{ exp}}$: experimental partial half-life calculated from the value of the transition probability of the quoted α transition;

$(T_{1/2})_{i \text{ th}}$: theoretical partial half-life.

All even-even nucleus transitions from ground state to ground state have, by definition, $F = 1$.

Partial half-lives $(T_{1/2})_{i \text{ th}}$ can be calculated by means of a selected theory of α decay [60Mi H], or derived from equation (3.13). In this case, the A and B constants for even-odd or odd-odd nuclei are derived by interpolation of the adjacent values of even-even nuclei.

For the even-even nucleus α transitions to excited levels, the hindrance factor which is caused, for the initial levels, by the angular momentum carried off by the α particle, slowly and steadily increases with the energy of the excited level (e.g. ^{226}Ra , $F \simeq 5$ for the second excited level).


Some transitions of even-odd and odd-odd nuclei, which generally do not go towards the ground state, exhibit a low hindrance factor. They are called "favored" (e.g. ^{241}Am) and

frequently correspond to a transition between initial and final states with identical spins and parities.

3.2.2 β^- decay

During a β^- decay, the mass number A remains unchanged and the atomic number Z varies by one unit.

β^- decay: an electron and an anti-neutrino are emitted, resulting from the transformation of a nucleus neutron to a proton.




The energy of a β^- transition, corresponding to the maximum β -energy emitted, and ignoring the recoil energy, is:

$$E_{\beta^-max} = Q^- - E_i$$

where Q^- : energy of disintegration, equal to the difference in atomic masses between the ground states of the parent and daughter, and E_i : energy of the level to which the decay occurs.

β^+ decay, the process is described similarly:



$$E_{\beta^+max} = Q^+ - 2m_0c^2 - E_i \quad (3.16)$$

where Q^+ : energy of disintegration, with

$$2m_0c^2 = 1,021\,998 \text{ MeV}$$

m_0 being the electron mass at rest.

A β^+ disintegration can occur, if:

$$Q^+ - E_i > 2m_0c^2 \quad (3.17)$$

Mean energy: the energy of a β transition is shared between the electron and the neutrino with a continuous distribution for the two particles extending from 0 up to $E_{\beta max}$.

If $N(E)dE$ is the number of β particles with energy between E and $E + dE$, the mean energy is written:

$$\bar{E} = \int_0^{E_{\beta max}} E N(E) dE \bigg/ \int_0^{E_{\beta max}} N(E) dE \quad (3.18)$$

Probability of β decay

The β decay is governed by the laws of weak interaction between the electron-neutrino pair and the nucleus and incorporates a very low coupling constant $g = 1,41 \times 10^{-56} \text{ J.cm}^3$.

The probability per unit time for an atomic nucleus to decay by a given β transition is [66Sc H, 66Ko E], in natural units ($\hbar = m_0 = c = 1$):

$$\lambda_n = \int_1^{W_0} N(W) dW \quad (3.19)$$

$$\lambda_n = \frac{g^2}{2\pi^3} \int_1^{W_0} W(W^2 - 1)^{1/2} (W_0 - W)^2 F(Z, W) L_0(Z, W) (1 + \delta_R(Z, W)) C_n dW \quad (3.20)$$

where:

n	:	order of forbiddenness of the transition
$N(W) dW$:	number of β particles emitted per unit time with energy between W and $W + dW$
$F(Z, W)$:	Fermi function
$L_0(Z, W)$:	screening correction
$\delta_R(Z, W)$:	radiative corrections
C_n	:	shape factor
Z	:	atomic number of the daughter
W	:	normalized total energy of the β particle $W = 1 + E/m_0$
W_0	:	maximum normalized total energy of the β particle $W_0 = 1 + E_{\beta\pm_{max}}/m_0$

The term $W(W^2 - 1)^{1/2} (W_0 - W)^2$ is the statistical factor determining the distribution of energy between the electron and the neutrino.

The Fermi function $F(Z, W)$ and the term $L_0(Z, W)$ take into account the perturbation of the electron wave function by the electric field of the nucleus. This Coulomb interaction causes at low spectral energies an enrichment of electrons and a depletion of positrons (Figure 3.1).




Figure 3.1: Examples of the typical β^- (left) and β^+ (right) spectra of an allowed transition.

The term $(1 + \delta_R(Z, W))$ takes into account the radiative corrections; for example the emission of photons due to internal bremsstrahlung during the decay.

The shape factor may be deduced from the transition characteristics (nuclear matrix elements, etc.).

Extensive compilation of experimental shape factors can be found in [76Be H].

Descriptions of the theoretical calculations for allowed and unique forbidden transitions can be found in [11Mo X].

Types of transitions and selection criteria

The initial and final nuclear states have well-defined total angular momenta with quantum numbers J_i and J_f . The electron-neutrino pair can carry off an orbital angular momentum with quantum number L_β , as well as an intrinsic spin S_β which can be: 1, -1 or 0. The conservation of the total angular momentum requires that:

$$J_i = J_f + L_\beta + S_\beta \quad (3.21)$$

$$\Delta J = J_i - J_f \quad (3.22)$$

as well as the parity relationship:

$$\pi_f \pi_i = (-1)^{L_\beta} \quad (3.23)$$

The transitions in which $L_\beta = 0$ are called allowed, those in which $L_\beta = n$ are called forbidden to the n^{th} order.

From this, selection criteria can be derived:

Classification of β -transition types

Type	Order of forbiddenness	ΔJ	$\pi_f \pi_i$
Super-allowed Allowed		0	+1
		0,+1	+1
Forbidden unique	1	± 2	-1
	2	± 3	+1
	3	± 4	-1
	4	± 5	+1
	$(\Delta J - 1)$ nth	> 1	$(-1)^{(\Delta J - 1)}$
Forbidden (non-unique)	1	0, ± 1	-1
	2	± 2	+1
	3	± 3	-1
	4	± 4	+1
	(ΔJ) nth	> 1	$(-1)^{\Delta J}$

For an allowed transition, the shape factor C_0 is independent of the energy and, in this case, the probability λ_0 of the transition is:

$$\lambda_0 = \frac{g^2}{2\pi^3} f_0(Z, W_0) C_0 \quad (3.24)$$

where:

$$f_0(Z, W_0) = \int_1^{W_0} W(W^2 - 1)^{1/2} (W_0 - W)^2 F(Z, W) dW \quad (3.25)$$

is the Fermi integral.

For unique forbidden transitions, the shape factor may be evaluated as a function of the energy. For non-unique forbidden transitions, it cannot be evaluated, because some matrix elements are unknown.

Comparative half-life, $\lg ft$

In order to obtain information on the shape factor, and on level spins and parities, and to make a rough classification of β transitions as a function of the forbiddenness degree, it is of great interest to calculate the comparative half-lives, ft , with:

$$t_i = \frac{T_{1/2}}{P_{\beta_i}} = \frac{\ln 2}{\lambda_i} \quad (3.26)$$

where:

- $T_{1/2}$: half-life in seconds,
- P_{β_i} : probability of the transition,
- λ_i and t_i : probability and “partial half-life” of the transition respectively.

For an allowed transition:

using (3.23) and (3.25)

$$f_0 t = \frac{2\pi^3 \ln 2}{g^2 C_0} \quad (3.27)$$

For forbidden transitions: [66Sc H]

$$f_n t = \frac{2\pi^3 \ln 2}{g^2 \eta^2} \quad (3.28)$$

with:

$$f_n = \int_1^{W_0} W(W^2 - 1)^{1/2} (W_0 - W)^2 F(Z, W) \frac{C_n}{\eta^2} dW \quad (3.29)$$

where η^2 is a combination of nuclear matrix elements.

The function f_n can only be calculated for unique transitions. However, for non-unique forbidden transitions, it is accepted that $C_n \simeq C_0$ to determine experimental values of ft .

In view of the extremely wide variety of the ft values, it is preferable to calculate and tabulate $\lg ft$ rather than ft [72Go N], as $\lg ft$ varies by about 3 for so-called super-allowed transitions, and increases with the order of forbiddenness up to a value above 20 for fourth-forbidden transitions.


β transitions are distributed as a function of $\lg ft$ as follows [73Ra S, 98Si B]:

Type of transition	$\lg ft$
Super-allowed	3,48–3,5
Allowed	3,5–12,5
First-forbidden	5,1–17
Second-forbidden	11–15,5

Type of transition	lg ft
Unique first-forbidden	8,5–11,7 (lg f_1t)
Unique second-forbidden	12,8–19
Forbiddenness greater than second	> 17,6

3.2.3 Electron capture decay

In this process, an orbital electron is captured by the nucleus:



A proton is transformed into a neutron and the atomic number decreases by one unit. The capture is accompanied by the emission of a monoenergetic neutrino, with energy q_x ($x = \text{K}, \text{L}, \text{M}, \text{etc.}$):

$$q_x = Q^+ - E_i - E_x \quad (3.30)$$

where:

- Q^+ : energy of disintegration corresponding to the difference in atomic masses between parent and daughter ground states,
- E_i : energy of the nuclear level towards which the electron capture occurs,
- E_x : binding energy of the electron in the X shell (or the subshell) in the daughter atom,
- $Q^+ - E_i$: transition energy.

The energy condition determining whether capture can occur in the X shell is:

$$Q^+ - E_i \geq E_x \quad (3.31)$$

furthermore, if:

$$Q^+ - E_i \gg E_x \quad (3.32)$$

the electron capture is far more probable in the K shell than in the other shells. Moreover, if:

$$Q^+ - E_i \geq 2 m_0 c^2 \quad (3.33)$$

the β^+ emission is in competition with the electron capture. For an electron capture transition, designated by ϵ , with a probability P_ϵ , the relative capture probabilities are:

$$P_K + P_L + P_M + P_N + \dots = 1 \quad (3.34)$$

Recoil energy: this is obtained from the laws of momentum and energy conservation as

$$E_r = \frac{q_x^2}{2 M_N c^2} \quad (3.35)$$

where M_N : mass of the nucleus.

It is generally ignored, except for ${}^7\text{Be}$, where $E_r = 57 \text{ eV}$.

Probability for a transition by electron capture

Electron capture is a weak interaction process to which the β -disintegration theory can be applied directly.

Allowed transitions: according to theory, the probability for capture per unit time in the K shell is given by the expression [60Bo R, 63Ba J]:

$$\lambda_k = \frac{g^2 |M_{0,1}|^2}{4\pi^2} q_K^2 g_K^2 B_K \quad (3.36)$$

The capture probabilities in the subshells L_1 and L_2 are given by identical expressions, in which q_K^2 is replaced by $q_{L_1}^2$ or $q_{L_2}^2$, g_K^2 by $g_{L_1}^2$ or $f_{L_2}^2$ and B_K by B_{L_1} or B_{L_2} . Capture in the L_3 subshell is negligible for allowed transitions.

$ M_{0,1} ^2$:	nuclear matrix element,
q_K, q_{L_1}, q_{L_2}	:	neutrino energies,
f, g	:	components of the electron radial wave functions,
B_K, B_{L_1}, B_{L_2}	:	exchange factors.

Except for the B_K factor, expression (3.36) is identical to that given for the β disintegration probability, the statistical factor being reduced to $\frac{\pi}{2} q_K^2 g_K^2$.

The exchange factor B , which takes into account atomic influences on the capture probability, incorporates two effects: the incomplete recovery of initial and final atomic states due to the charge change in the nucleus, and the exchange effect itself, since the capture in a shell can be achieved by several processes which are not experimentally distinguishable. For instance, a L_1 vacancy can be obtained mainly by direct capture of an L_1 electron, but also by K capture with simultaneous exchange of L_1 and K electrons, or by capture of an M_1 electron with simultaneous exchange of M_1 and L_1 electrons.

These two effects, recovery and exchange, have opposite signs and partially counteract each other. The recovery effect predominates for low atomic numbers, and the exchange effect for high atomic numbers.

The capture probabilities P_K, P_L, P_{L_1}, \dots are normally calculated by means of the ratios:

$$\frac{P_L}{P_K} = \frac{P_{L_1}}{P_K} \left[1 + \frac{P_{L_2}}{P_{L_1}} \right] \quad (3.37)$$

$$\frac{P_{L_1}}{P_K} = \frac{g_{L_1}^2 q_{L_1}^2}{g_K^2 q_K^2} X^{L_1/K} \quad \text{with} \quad X^{L_1/K} = \frac{B_{L_1}}{B_K} \quad (3.38)$$

$$\frac{P_{L_2}}{P_{L_1}} = \frac{f_{L_2}^2 q_{L_2}^2}{g_{L_1}^2 q_{L_1}^2} \frac{B_{L_2}}{B_{L_1}} \simeq \frac{f_{L_2}^2}{g_{L_1}^2} \frac{B_{L_2}}{B_{L_1}} \quad (3.39)$$

The ratios $\frac{g_{L_1}^2}{g_K^2}$ and $\frac{f_{L_2}^2}{g_{L_1}^2}$ have been calculated by several authors [69Su Y, 69Be H, 70Ma M] together with exchange factors [70Su Y, 70Fa J, 70Va E].

The capture probabilities in the M shell are determined similarly to those in the K and L shells by means of the ratios:

$$\frac{P_M}{P_L} = \frac{P_M}{P_{L_1}} \frac{P_{L_1}}{P_L} \quad (3.40)$$

where $\frac{P_{L_1}}{P_L}$ is deduced from (3.37) and (3.39), moreover:

$$\frac{P_M}{P_{L_1}} = \frac{P_{M_1}}{P_{L_1}} \left[1 + \frac{P_{M_2}}{P_{M_1}} \right] \quad (3.41)$$

$$\frac{P_M}{P_{L_1}} \simeq \frac{g_{M_1}^2 q_{M_1}^2}{g_{L_1}^2 q_{L_1}^2} \left[1 + \frac{f_{M_2}^2}{g_{M_1}^2} \right] X^{M_1/L_1} \quad (3.42)$$

with $X^{M_1/L_1} = \frac{B_{M_1}}{B_{L_1}}$.

Captures in the M_3 , M_4 , M_5 subshells are assumed to be negligible.

More general expressions including captures in the lower shells may be written [70Ma M]:

$$\frac{P_{MN}}{P_L} = \frac{P_{MN}}{P_{L_1}} \frac{P_{L_1}}{P_L} \quad (3.43)$$

$$\frac{P_{MN}}{P_{L_1}} = \frac{P_{M_1}}{P_{L_1}} \left[1 + \frac{P_{M_2} + P_{N_1} + P_{N_2} + \dots}{P_{M_1}} \right] \quad (3.44)$$

Unique forbidden transitions: the capture probability in the K shell is written [60Bo R, 70Ma M]:

$$\lambda_K = \frac{g^2 |M_{\Delta J}|^2 (q_K r_0)^{2(\Delta J-1)}}{4\pi^2 [(2\Delta J-1)!!]^2} q_K^2 g_K^2 B_K \quad (3.45)$$

where r_0 : nucleus radius in units of \hbar/m_0c .

$$n = \Delta J - 1 \quad \Delta J = 2, 3, \dots \quad \pi_f/\pi_i = (-1)^n \quad (3.46)$$

The capture probabilities in the L_1 and L_2 subshells are identically obtained and the contribution due to the capture in the L_3 subshell must be added. They are determined by means of the following ratios:

$$\frac{P_L}{P_K} = \frac{P_{L_1}}{P_K} \left[1 + \frac{P_{L_2}}{P_{L_1}} + \frac{P_{L_3}}{P_{L_1}} \right] \quad (3.47)$$

$$\frac{P_{L_3}}{P_{L_1}} = \frac{3(\Delta J-1)(2\Delta J-1)}{(q_{L_1} r_0)^2} \frac{g_{L_3}^2 B_{L_3}}{g_{L_1}^2 B_{L_1}} \quad (3.48)$$

$$\frac{P_L}{P_K} \simeq \frac{g_{L_1}^2 q_{L_1}^{2\Delta J}}{g_K^2 q_K^{2\Delta J}} X^{L_1/K} \left[1 + \frac{f_{L_2}^2 B_{L_2}}{g_{L_1}^2 B_{L_1}} + \frac{3(\Delta J-1)(2\Delta J-1)}{(q_{L_1} r_0)^2} \frac{g_{L_3}^2 B_{L_3}}{g_{L_1}^2 B_{L_1}} \right] \quad (3.49)$$

lg *ft*: the comparative half-lives for electron capture transitions are calculated similarly to those of β transitions [72Go N].

3.2.4 ϵ/β^+ ratio

For nuclides decaying by electron capture and β^+ disintegration, when the electron capture is in competition with the β^+ emission, the ratio of probabilities $P\epsilon/P_{\beta^+}$, or more simply ϵ/β^+ , is written:

$$\frac{\epsilon}{\beta^+} = \frac{\epsilon}{\beta^+} [P_K + P_L + P_M + \dots] \quad (3.50)$$

$$\frac{\epsilon}{\beta^+} = \frac{\epsilon P_K}{\beta^+} \left[1 + \frac{P_L}{P_K} + \frac{P_M}{P_K} + \dots \right] \quad (3.51)$$

The terms in square brackets may be calculated by equations (3.37) to (3.49).

Let $\epsilon P_K = \epsilon_K$; $\frac{\epsilon_K}{\beta^+}$ is given by:

$$\frac{\epsilon_K}{\beta^+} = \frac{\pi}{2} \frac{q_K^2 g_K^2 B_K}{f_0} \quad (3.52)$$

for an allowed transition;

$$\frac{\epsilon_K}{\beta^+} = \frac{\pi}{2} \frac{q_K^{2\Delta J} g_K^2 B_K}{f_{\Delta J-1}} \quad (3.53)$$

for a unique forbidden transition.

3.3 γ transitions

A γ transition occurs when a nucleus is de-excited towards a lower energy level that may or may not, be the ground state. This leads to the emission of a γ photon or to the ejection of an electron from the electronic shells, which is called a conversion electron, and very rarely, when energy conditions make it possible, of an electron-positron pair.

The γ transition probability is:

$$P_g = I_\gamma + I_{ce} + I_{e^\pm} \quad (3.54)$$

where I_γ , I_{ce} and I_{e^\pm} are the γ emission, conversion electron emission and electron-positron pair emission intensities, respectively.

3.3.1 γ photon emission

The energy of the emitted γ photon is:

$$E_\gamma = (E_i - E_f) - E_r \quad (3.55)$$

where $E_i - E_f$: energy difference between the initial and the final level of the γ transition, E_r : recoil energy of the nucleus in its final state, given by:

$$E_r = \frac{(E_\gamma)^2}{2M_N c^2} \quad (3.56)$$

M_N : mass of the recoiling daughter nucleus.

The recoil energy is negligible, except for high γ energies and low atomic numbers. For example, for the 1369 keV γ emission of ^{24}Na , $E_r = 42$ eV.

3.3.2 Internal conversion

The de-excitation energy of the nucleus can also be transferred directly to an X-shell electron ($X = K, L, M \dots$) which is ejected from the atom, carrying off the energy:

$$E_{ce_X} = E_\gamma - E_X \quad (3.57)$$

where E_X : binding energy of the electron in the X shell.

Internal conversion coefficients:

For a given transition, the internal conversion coefficient of the electron in the K shell is defined by:

$$\alpha_K = \frac{I_{ce_K}}{I_\gamma} \quad (3.58)$$

where I_{ce_K} and I_γ are the K conversion electron emission and γ emission intensities, respectively.

The following quantities are similarly defined:

$$\alpha_L = \frac{I_{ce_L}}{I_\gamma}, \quad \alpha_{L_i} = \frac{I_{ce_{L_i}}}{I_\gamma} \quad (i = 1, 2, 3) \quad (3.59)$$

with $\alpha_L = \alpha_{L_1} + \alpha_{L_2} + \alpha_{L_3}$

$$\alpha_{M_i} = \frac{I_{ce_{M_i}}}{I_\gamma} \quad (i = 1, 2, 3, 4, 5) \quad (3.60)$$

The total conversion coefficient is:

$$\alpha_t = \alpha_K + \alpha_L + \alpha_M + \dots = \frac{I_{ce}}{I_\gamma} \quad (3.61)$$

I_{ce} is the total conversion electron emission intensity for the related transition.

This leads to the following relations, neglecting P_{e^\pm} :

$$I_{ce} = \frac{\alpha_t}{1 + \alpha_t} P_g, \quad I_\gamma = \frac{P_g}{1 + \alpha_t}, \quad I_{ce_X} = \frac{\alpha_X}{1 + \alpha_t} P_g \quad (X = K, L, M \dots) \quad (3.62)$$

The K/L , K/LM , $K/LM\dots$, ratios are also employed and are defined as follows:

$$\frac{K}{L} = \frac{I_{ce_K}}{I_{ce_L}} = \frac{\alpha_K}{\alpha_L} \quad (3.63)$$

$$\frac{K}{LM} = \frac{I_{ce_K}}{I_{ce_L} + I_{ce_M}} = \frac{\alpha_K}{\alpha_L + \alpha_M} \quad (3.64)$$

$$\frac{K}{LM\dots} = \frac{I_{ce_K}}{I_{ce_L} + I_{ce_M} + \dots} = \frac{\alpha_K}{\alpha_L + \alpha_M + \dots} \quad (3.65)$$

Internal conversion depends on the initial state of the electron (atomic shell or subshell), on the atomic number Z and on the nuclear transition multipolarity. It increases with multipolarity and decreases with increasing γ energy, becoming negligible for high energies.

Internal conversion is influenced by the finite size of the nucleus in relation with two processes: a static one owing to the finite distribution of nuclear charge, and a dynamic one (penetration process) which expresses the non-zero conversion probability within the nuclear volume. This penetration effect can become significant only in some specific cases such as: γ transition with high hindrance, heavy nuclei, etc.

3.3.3 Internal pair creation

This process is possible only if the transition energy is greater than $2 m_0 c^2 = 1,022$ MeV, with the electron-positron pair carrying off the energy:

$$E (e^\pm) = E_\gamma - 2 m_0 c^2 \quad (3.66)$$

in accordance with a distribution extending from 0 to $E (e^\pm)$ for each of the two particles (e.g. decay of the first excited level of ^{22}Na).

The internal pair creation coefficient is defined by:

$$\alpha_\pi = \frac{N_{e^\pm}}{N_\gamma} \quad (3.67)$$

where N_{e^\pm} is the number of created electron-positron pairs and N_γ the number of emitted photons. The internal pair creation coefficient increases with increase of γ transition energy [68Wi R].

α_π is typically of the order of $10^{-4} - 10^{-3}$.

3.3.4 Multipolarity

Quantum theory permits a classification of γ transitions into multiple order transitions. This order is a function of the orbital angular momentum, of quantum number L , carried off by the photon during the transition. If $L = 0$, the transition is called monopole, $L = 1$ dipole, $L = 2$ quadrupole, etc.

If J_i and J_f are the total angular momenta quantum numbers of the initial and final levels of the γ transition, the vectorial relationship between the angular momenta imposes the relation:

$$|J_i - J_f| \leq L \leq |J_i + J_f| \quad (3.68)$$

Because of the strong dependence of the γ transition probability with L , the γ transition probability decreases rapidly when L increases.

More often, $L = |J_i - J_f|$ can be in competition with $L = |J_i - J_f| + 1$ and, more rarely with $L = |J_i - J_f| + 2$. Moreover, the angular momentum carried off by the photon cannot be zero. Consequently, a $0 \rightarrow 0$ transition cannot occur except by internal conversion or by internal pair creation.

Furthermore, γ radiation is divided into “electric” and “magnetic” radiations.

For “electric” radiation, i.e. emitted by the oscillation of electrical charges, the parity change is $(-1)^L$.

For “magnetic” radiation, caused by the magnetic moment of the nucleus, the parity change is $(-1)^{(L+1)}$.

A γ transition can be a mixture of two (or more rarely three) multipole transitions. In order to conserve parity, two transitions in competition will have the multiple order 2^L and 2^{L+1} , one electric and the other one magnetic, or inversely (e.g. mixture of M1 + E2).

Expression (3.68) implies that a transition with:

- $J_i = J_f = 1/2$ is necessarily pure E1 or M1 ;
- J_i or $J_f = 0$, L has only one possible value, *i.e.* the multipole character of the transition is pure E_L or M_L . For example, generally even-even nuclei have a $J_f = 0$ ground state level, then the occurring transition has a pure multipolarity character.

Note also that for a $J_f = J_i$ transition without parity change, a monopole transition E0 can be in competition with the mixture M1 + E2.

The classification of γ transitions and their possible mixtures is established as follows:

$L = J_i - J_f $	$\pi_f/\pi_i = -1$		$\pi_f/\pi_i = +1$	
	Transition	Possible mixture	Transition	Possible mixture
0 (0 \rightarrow 0)	Forbidden		Forbidden	
0 (1/2 \rightarrow 1/2)	E1		M1	
0 (1 \rightarrow 1)	E1	M2	M1	E2
1	E1	M2	M1	E2
2	M2	E3	E2	M3
3	E3	M4	M3	E4
4	M4	E5	E4	M5
even L (J_i or $J_f = 0$)	ML		EL	
odd L (J_i or $J_f = 0$)	EL		ML	

The mixture is characterized, e.g. $y\%M1 + x\%E2$, by $\delta^2 = E2/M1$ which is the ratio of the probability of the E2 transition to that of the M1 transition [53Bi L].

Then, $x = 100 \delta^2 / (1 + \delta^2)$ and $y = 100 / (1 + \delta^2)$

3.3.5 Mean life times of excited levels

The mean life times of γ transitions are a function of the energy, the atomic number and the multipolarity. They range from 10^{-14} second to several years for certain M4 and E5 low energy transitions.

On the basis of the simple theory of the shell model, Weisskopf [51We V] calculated the probability λ per unit time (transition rate) for each type of transition. The mean life time $\tau = 1/\lambda$ decreases with $(E_\gamma)^{2L+1}$.

Moreover, for the same energy and atomic number, and the same multipole order, we have the relationship:

$$\frac{\tau_{elec}}{\tau_{magn}} \simeq 2 \times A^{2/3} \quad (3.69)$$

In fact, owing to the simplicity of the adopted model, the experimental values deviate substantially from calculated values in most cases. However, in view of the cumulative results [66Go N], it may be considered that if:

$(T_{1/2})_{exp}/[(T_{1/2})_{cal}]_{M1}$ (M1) < 0,1	the transition is not M1
$(T_{1/2})_{exp}/[(T_{1/2})_{cal}]_{M2}$ (M2) < 1,0	the transition is not M2
$(T_{1/2})_{exp}/[(T_{1/2})_{cal}]_{M3}$ (M3) < 0,1	the transition is not M3
$(T_{1/2})_{exp}/[(T_{1/2})_{cal}]_{M4}$ (M4) < 0,033	the transition is not M4
$(T_{1/2})_{exp}/[(T_{1/2})_{cal}]_{E1}$ (E1) < 100	the transition is not E1
$(T_{1/2})_{exp}/[(T_{1/2})_{cal}]_{E2}$ (E2) < 0,001	the transition is not E2
$(T_{1/2})_{exp}/[(T_{1/2})_{cal}]_{E3}$ (E3) < 0,01	the transition is not E3

3.4 Secondary phenomena accompanying nuclear transformations

Transitions are always accompanied by secondary effects of low probability which can be theoretically predicted and calculated, and which appear as emission of electrons and photons. These secondary effects are frequently masked by the main radiations or by effects induced by radiations (e.g. Compton effect), making them difficult to measure.

3.4.1 Internal bremsstrahlung accompanying β disintegration

After its emission, an electron can be slowed down in the field of the nucleus, and the bremsstrahlung energy appears in the form of a photon [68Pe B]. (This internal bremsstrahlung must be distinguished from the external bremsstrahlung occurring in the absorbent medium). The transition energy is distributed between the electron, the neutrino and the internal bremsstrahlung photon:

$$E_{\beta max} = E_{\beta} + E_{\nu} + E_{ph} \quad (3.70)$$

The spectral distribution reveals a significant decrease in bremsstrahlung photons with their energy. The ratio of the emission probabilities for photons and β particles is expressed by:

$$\frac{P_{ph}}{P_{\beta-}} = \frac{4}{3} \frac{\alpha}{\pi} \frac{1}{\lg(2 W_0 - \zeta)} \quad (3.71)$$

where:

- α = 1/137,035 999 074 (44), fine structure constant [10Mo P]
- W_0 : maximum energy in $m_0 c^2$ units
- ζ : constant between 2,20 and 2,35 depending on the transition type

For ^{35}S and ^{32}P , this ratio is 2×10^{-5} and 2×10^{-3} respectively. Theoretical and experimental values agree to within 20 %.

3.4.2 Internal bremsstrahlung accompanying electron capture

The origin of this photon radiation is the same as that accompanying β disintegration [68Zy J]. The available energy $E_{\beta max} = Q - E_K$ for a capture in the K shell is shared between the

neutrino and the bremsstrahlung photon.

As there are no external bremsstrahlung, this process is fairly easy to analyze, and theoretical and experimental results agree within less than 15 %.

The analysis of the spectrum is of great interest as it permits direct determination of the transition energy Q . The maximum energy $E_{\beta max}$, is obtained from a plot [40Mo P]:

$$\left(\frac{N(E)}{E}\right)^{1/2} = C (E_{\beta max} - E) \quad (3.72)$$

where $N(E)$ is the number of bremsstrahlung photons with energy E and C is a constant.

The ratio of the number of emitted internal bremsstrahlung quanta and the number of K-capture events is given by [73Sp A]:

$$\frac{N_{ph}}{N_K} = 2 \times 10^{-4} \left(\frac{E_{\beta max}}{m_0 c^2}\right)^2 f(Z) \quad (3.73)$$

where $f(Z)$ decreases from 0,9 to 0,3 when Z increases from 4 to 68.

For ^{55}Fe , this ratio is 3×10^{-5} .

3.4.3 Internal ionization and excitation accompanying β decay

This process is attributed to the sudden change in the nuclear charge during β decay [72Sh S]. In internal ionization, it appears by the simultaneous emission of a β particle, a neutrino and an atomic shell electron, with the following energy relationship:

$$E_{\beta max} = E_{\beta} + E_{\nu} + E_e + E_X \quad (3.74)$$

where:

- $E_{\beta max}$: maximum β -transition energy
- E_{β} , E_{ν} , E_e : energies of the three emitted particles
- E_X : binding energy of the shell (or subshell) from which the electron is emitted

The internal excitation is produced by the electron transition to an unoccupied state of a less bound shell.

The ionization and internal excitation in the K shell are best known. They are accompanied by the creation of a vacancy; the resulting emission of a X_K photon or a K-Auger electron makes the phenomenon measurable.

The internal ionization probability in the K shell is roughly proportional to Z^{-2} , going from 3×10^{-3} for ^{32}P to 1×10^{-4} for ^{204}Tl . Theoretical and experimental results agree within about 15 %.

3.4.4 Internal ionization and excitation accompanying electron capture

When an atom disintegrates by electron capture in the K shell, a double vacancy can be created [53Pr H, 69St P]. The second electron can be ejected from the atom (internal ionization or shake-off) or can undergo a transition to an unoccupied state of a less bound shell (internal excitation or shake-in). The process occurs before the rearrangement of the orbital electrons. In the case of internal ionization, the transition energy, decreased by the binding energy, is shared between the neutrino and the ejected electron.

The probability P_{KK} for the formation of a double K-shell vacancy is of the order of $0,18 Z^{-2}$, while that for the electron ejection is of the order of $0,08 Z^{-2}$.

In the case of ^{55}Fe , the total probability for double vacancies in the K shell was measured to be $1,3 \times 10^{-4}$ [91Ca J].

3.4.5 Other processes

It is also worth mentioning the secondary phenomena of very low probability occurring during a γ transition [68Le H] such as the creation of monoenergetic positron, double photon emission, double γ -e emission, double internal conversion.

3.5 Electron rearrangement

Electron capture and internal conversion (as well as the ionization and internal excitation which can be ignored in the following considerations) are processes giving rise to vacancies in the electron cloud. The filling of a vacancy is accompanied by the emission of an X-ray or an Auger electron, creating new vacancies in the less bound shells. The initial vacancy is thus transferred by cascade to the peripheral shell. The energy liberated corresponds to the binding energy of the electron in the shell in which the initial vacancy was produced.

3.5.1 K shell

Vacancies in the K shell

The probability for K vacancies created during an electron capture transition with probability P_e , is $P_e P_K$ (P_K : probability for the formation of a vacancy in the K shell).

The probability for K vacancies created during an electron conversion process of a γ transition with the probability P_g is: $P_g \frac{\alpha_K}{1 + \alpha_t}$.

For a given radionuclide, the total number of K vacancies created, following one disintegration of the initial nuclide, and taking account of all transitions, is:

$$n_K = \sum P_e P_K + \sum P_g \frac{\alpha_K}{1 + \alpha_t} \quad (3.75)$$

K-shell fluorescence yield

The K shell has no subshells. The K-shell fluorescence yield ω_K is the probability that the filling of a vacancy in the K shell is accompanied by a X_K radiation, $0 \leq \omega_K \leq 1$.

$$\omega_K = \frac{n_{X_K}}{n_{X_K} + n_{A_K}} = \frac{I_{X_K}}{I_{X_K} + I_{A_K}} \quad (3.76)$$

where:

n_{X_K} : number of X_K photons

n_{A_K} : number of K-Auger electrons

I_{X_K} : emission intensity for X_K photons

I_{A_K} : emission intensity for K-Auger electrons

The K-Auger yield is derived from the fluorescence yield:

$$a_K = 1 - \omega_K \quad (3.77)$$

As well as the number of X_K photons and Auger electrons is:

$$I_{X_K} = \omega_K(I_{X_K} + I_{A_K}) = \omega_K n_K \quad (3.78)$$

$$I_{A_K} = a_K(I_{X_K} + I_{A_K}) = a_K n_K \quad (3.79)$$

The fluorescence yield ω_K depends exclusively on the atomic number Z , except for low values of Z , where it may be influenced by the chemical state.

$\omega_K \ll 1$ for low atomic numbers. It approaches 1 for high atomic numbers.

K X-rays

The K X-ray energy emitted for a K-X transition (X= L, M, N ...) is written as:

$$E_{X_K} = E_K - E_X \quad (3.80)$$

where E_K and E_X are the binding energies of the electrons in the K and X shells respectively.

Consequently, the energy of the K X-rays is perfectly defined. They are usually classified by series, consisting of the following rays:

$$K\alpha \quad \left\{ \begin{array}{ll} K\alpha_2 & K-L_2 \\ K\alpha_1 & K-L_3 \end{array} \right.$$

$$K'\beta_1 \quad \left\{ \begin{array}{ll} K\beta_3 & K-M_2 \\ K\beta_1 & K-M_3 \\ K\beta_5'' & K-M_4 \\ K\beta_5 & K-M_5 \end{array} \right.$$

$$K'\beta_2 \quad \left\{ \begin{array}{ll} K\beta_2 \quad \left\{ \begin{array}{ll} K\beta_2'' & K-N_2 \\ K\beta_1' & K-N_3 \end{array} \right. \\ K\beta_4 \quad \left\{ \begin{array}{ll} K\beta_4'' & K-N_4 \\ K\beta_4' & K-N_5 \end{array} \right. \end{array} \right.$$

Transitions KL_1 , KM_1 , KN_1 are forbidden.

For $Z > 49$, it is necessary to take the KO_2 and KO_3 rays into account.

The emission probability for specific groups of K X-rays can be calculated from the total K X-ray emission probability, P_{X_K} , and from the emission probability ratios:

$$I(K_\alpha) = I_{X_K} \left[1 + \frac{P(K_\beta)}{P(K_\alpha)} \right]^{-1}$$

$$I(K_\beta) = I_{X_K} \left[1 + \frac{P(K_\alpha)}{P(K_\beta)} \right]^{-1}$$

$$I(K_{\alpha_1}) = I(K_\alpha) \left[1 + \frac{P(K_{\alpha_2})}{P(K_{\alpha_1})} \right]^{-1}$$

$$I(K_{\alpha_2}) = I(K_\alpha) \left[1 + \frac{P(K_{\alpha_1})}{P(K_{\alpha_2})} \right]^{-1}$$

The emission probability ratios $P(K_\beta)/P(K_\alpha)$ and $P(K_{\alpha_2})/P(K_{\alpha_1})$ have been measured by several authors [95Sc E]. They have also been calculated theoretically [74Sc J].

K-Auger electrons

A vacancy in the K shell is filled by an electron coming from a less bound shell. The available energy is transferred to an electron of another shell, which is also less bound. The latter electron is then ejected (Auger electron).

The energy of the K-Auger electron is:

$$E_{AK} = E_K - E_X - E_Y - \Delta E \quad (E_Y \leq E_X) \quad (3.81)$$

where E_K , E_X , E_Y , are the binding energies of the electrons in the K-shell and in the two less bound shells (X, Y) (or subshells). ΔE is a corrective term accounting for the fact that the binding energy of the excited atom is greater than that of the atom in its ground state.

Three groups of K-Auger electrons are distinguished. Using the conventional notation we have:

- KLL-Auger electrons with six components: KL_1L_1 , KL_1L_2 , KL_1L_3 , KL_2L_2 , KL_2L_3 , KL_3L_3 .
- K LX-Auger electrons (X = M, N...) with the following components : KL_1M_1 , KL_1M_2 , KL_1M_3 , $KL_1M_{4,5}$, KL_2M_1 , $KL_2M_{2,3}$, $KL_2M_{4,5}$, KL_3M_1 , $KL_3M_{4,5}$, KL_1N , KL_2N , KL_3N .
- KXY-Auger electrons (X = M, N...; Y = M, N...). The KXY-Auger electron spectrum is highly structured and the components are not be listed here.

As the KLL-Auger electrons are the most intense, the emission intensity P_{AK} is often written in the following form, with $\frac{I_{AKLX}}{I_{AKLL}} = \frac{K LX}{K LL}$:

$$I_{AK} = KLL \left(1 + \frac{K LX}{K LL} + \frac{K XY}{K LL} \right) \quad (3.82)$$

3.5.2 L shell

Vacancies created in the L shell

The vacancies created in the L shell can have several origins:

- electron capture in the L shell. For an electron capture transition with a probability P_ϵ , the vacancy number is: $P_\epsilon P_L$;
- internal conversion in the L shell. For a γ transition with a probability $P_g = I_\gamma + I_{ce}$, the vacancy number is: $P_g \alpha_L / (1 + \alpha_L)$;
- rearrangement following a K vacancy giving rise to a $K \rightarrow L$ vacancy transfer.

For a given radionuclide, the total number of L vacancies created per disintegration is written as follows, accounting for all transitions by electron capture, all γ transitions, and all $K \rightarrow L$ vacancy transfers:

$$n_L = n_{L_1} + n_{L_2} + n_{L_3} = \sum P_\epsilon P_L + \sum P_g \frac{\alpha_L}{1 + \alpha_t} + n_K n_{KL} \quad (3.83)$$

where:

- n_K : is the total number of K vacancies calculated by equation (3.75)
- n_{KL} : is the total number of L vacancies created by $K \rightarrow L$ vacancy transfers when one K vacancy is filled.

This vacancy transfer arises from the X_K emission ($K\alpha_1$ and $K\alpha_2$ rays) and from the emission of K-Auger electrons (KLL and KLX), with $I_{X_{K\alpha}} = K\alpha$, $I(\text{KLL}) = \text{KLL}$, etc.

$$n_{KL} = \omega_K \frac{K\alpha}{K\alpha + K\beta} + (1 - \omega_K) \frac{2\text{KLL} + \text{KLX}}{\text{KLL} + \text{KLX} + \text{KXY}} \quad (3.84)$$

it may also be written as:

$$n_{KL} = \omega_K \frac{1}{1 + \frac{K\beta}{K\alpha}} + (1 - \omega_K) \frac{2 + \frac{\text{KLX}}{\text{KLL}}}{1 + \frac{\text{KLX}}{\text{KLL}} + \frac{\text{KLY}}{\text{KLL}}} \quad (3.85)$$

L-shell fluorescence yields

For a L_i subshell ($i = 1, 2, 3$), this is the probability that the filling of a L_i vacancy is accompanied by the emission of a L_i X-ray.

This leads to the definition of three L-fluorescence yields ω_{L1} , ω_{L2} , ω_{L3} :

$$\omega_{Li} = \frac{I_{X_{Li}}}{n_{Li}} \quad (i = 1, 2, 3) \quad (3.86)$$

ω_{L1} , ω_{L2} and ω_{L3} , like ω_K , depend exclusively on the atomic number Z .

n_{Li} is the number of vacancies created in the L_i subshell for one disintegration of the parent nuclide, and $I_{X_{Li}}$ is the emission intensity for X_{Li} quanta.

L X-rays

The filling of a vacancy in the L shell by an electron coming from a less bound shell X ($X = M, N, \dots$) can be accompanied by the emission of X_L radiation of energy $E_{X_L} = E_L - E_X$, E_L and E_X being the binding energies of the L and X shells (or subshells).

The X_L series consists of the following main rays:

$$\begin{array}{l} L_\ell \{ \ell \quad L_3 - M_1 \\ L_\alpha \left\{ \begin{array}{l} \alpha_2 \quad L_3 - M_4 \\ \alpha_1 \quad L_3 - M_5 \end{array} \right. \\ L_\eta \{ \eta \quad L_2 - M_1 \end{array}$$

$$\begin{array}{l}
L_{\beta} \left\{ \begin{array}{l}
\beta_6 \quad L_3 - N_1 \\
\beta_4 \quad L_1 - M_2 \\
\beta_3 \quad L_1 - M_3 \\
\beta_{10} \quad L_1 - M_4 \\
\beta_9 \quad L_1 - M_5 \\
\beta_1 \quad L_2 - M_4 \\
\beta_2 \quad L_3 - N_5 \\
\beta_{15} \quad L_3 - N_4 \\
\beta_7 \quad L_3 - O_1 \\
\beta_5 \quad L_3 - O_{4, 5}
\end{array} \right. \\
\\
L_{\gamma} \left\{ \begin{array}{l}
\gamma_2 \quad L_1 - N_2 \\
\gamma_3 \quad L_1 - N_3 \\
\gamma_4 \quad L_1 - O_{2, 3} \\
\gamma_5 \quad L_2 - N_1 \\
\gamma_1 \quad L_2 - N_4 \\
\gamma_6 \quad L_2 - O_4
\end{array} \right.
\end{array}$$

L-Auger electrons

When a vacancy in the L shell is filled by an electron coming from a less bound shell (M, N, ...), the available energy is transferred to an electron of another less bound shell (M, N, ...). This latter electron is then ejected, it is called L-Auger electron.

The Auger yield a_{L_i} is the probability that a vacancy in the i^{th} shell is filled with an electron from a higher shell and is accompanied by the ejection of another less-bound electron. The L-Auger electron spectrum includes a large number of lines, which are practically indistinguishable.

Coster-Kronig transitions

A Coster-Kronig transition is a special type of Auger effect, it occurs when an L_i vacancy ($i = 1, 2$) moves from a subshell towards another less bound subshell, $L_1 \rightarrow L_2$, $L_1 \rightarrow L_3$, $L_2 \rightarrow L_3$ [65Fi R]. The ratio of the number of vacancies filled by Coster-Kronig transitions to the total number of initial vacancies in the given subshell, respectively f_{12} , f_{13} and f_{23} , are called Coster-Kronig yields.

A transition of this type is accompanied by the emission of an $L_i L_j X$ - Auger electron emitted from a less bound shell ($X = M, N \dots$). It carries off the energy:

$$E_{CK} \simeq E_{L_i} - E_{L_j} - E_X \quad (i = 1, 2 \quad j = 2, 3) \quad (3.87)$$

This emission, which occurs before the L_i subshell is filled, has the effect of increasing the number of L vacancies.

The new vacancies V_{L_1} , V_{L_2} and V_{L_3} are such that:

$$V_{L_1} + V_{L_2} + V_{L_3} > N_{L_1} + N_{L_2} + N_{L_3} \quad (3.88)$$

where N_{L_i} is the number of primary vacancies. The V_{L_i} are calculated by the following equations:

$$V_{L_1} = N_{L_1} \quad (3.89)$$

$$V_{L_2} = N_{L_2} + f_{12}N_{L_1} \quad (3.90)$$

$$V_{L_3} = N_{L_3} + f_{13}N_{L_1} + f_{23}(N_{L_2} + f_{12}N_{L_1}) \quad (3.91)$$

These expressions require that the initial vacancy distribution is given relatively to the total number of primary vacancies in the L shell:

$$N_L = N_{L_1} + N_{L_2} + N_{L_3} = 1 \quad (3.92)$$

hence:

$$V_{L_1} + V_{L_2} + V_{L_3} > 1 \quad (3.93)$$

Mean L-fluorescence yield

The mean fluorescence yield $\bar{\omega}_L$ is derived from the three individual fluorescence yields ω_{L_1} , ω_{L_2} and ω_{L_3} , and from the distribution of vacancies modified by the Coster-Kronig transitions.

$$\bar{\omega}_L = V_{L_1}\omega_{L_1} + V_{L_2}\omega_{L_2} + V_{L_3}\omega_{L_3} \quad (3.94)$$

$\bar{\omega}_L$ can also be expressed as a function of primary vacancies:

$$\bar{\omega}_L = N_{L_1}[\omega_{L_1} + f_{12}\omega_{L_2} + (f_{13} + f_{12}f_{23})\omega_{L_3}] + N_{L_2}(\omega_{L_2} + f_{23}\omega_{L_3}) + N_{L_3}\omega_{L_3} \quad (3.95)$$

The L X-rays probability is given by:

$$P_{X_L} = n_L\bar{\omega}_L \quad (3.96)$$

with n_L : total number of L-shell vacancies per disintegration.

Remark: While the yields ω_{L_1} , ω_{L_2} , ω_{L_3} , f_{12} , f_{13} and f_{23} depend exclusively on the atomic number, this is not the case for $\bar{\omega}_L$ which depends on the initial vacancies distribution, itself a function of the transitions which gave rise to it. Hence $\bar{\omega}_L$ is not constant for a given atomic number and must be measured or calculated for each radionuclide taking into account the characteristics and constants of the decay scheme.

Mean L-Auger yield

The mean L-Auger yield is derived from $\bar{\omega}_L$ for a given initial vacancy distribution, by the equation:

$$\bar{\omega}_L + \bar{a}_L = 1 \quad (3.97)$$

Where \bar{a}_L is composed by, taking into account the Coster-Kronig transitions:

$$\begin{aligned} \omega_{L_3} + a_{L_3} &= 1 \\ \omega_{L_2} + a_{L_2} + f_{23} &= 1 \\ \omega_{L_1} + a_{L_1} + f_{12} + f_{13} &= 1 \end{aligned}$$

And the total emission probability of the L-Auger electrons is:

$$P_{A_L} = n_L\bar{a}_L \quad (3.98)$$

With:

$$\begin{aligned}
 P_{A_L} &= P_{A_{L1}} + P_{A_{L2}} + P_{A_{L3}} \\
 P_{A_{L1}} &= a_{L1} N_{L1} = (1 - \omega_{L1} - f_{12} - f_{13}) N_{L1} \\
 P_{A_{L2}} &= a_{L2} (N_{L2} + f_{12} N_{L1}) = (1 - \omega_{L2} - f_{23}) (N_{L2} + f_{12} N_{L1}) \\
 P_{A_{L3}} &= a_{L3} (N_{L3} + f_{13} N_{L1} + f_{23} (N_{L2} + f_{12} N_{L1})) \\
 &= (1 - \omega_{L3}) ((N_{L3} + f_{13} N_{L1} + f_{23} (N_{L2} + f_{12} N_{L1})))
 \end{aligned}$$

Where $P_{A_{Li}}$ are the number of L_i Auger electrons.

Radiations accompanying a gamma transition

As an example of the effects of secondary phenomena accompanying the nuclear transformations, the expressions for the emission intensities of gamma rays, conversion electrons, Auger electrons and X-ray quanta are given below for a gamma transition with the transition probability P_g and the conversion coefficients α_K , α_L , α_M and α_t . If the vacancy transfer from the K to the L and from the L to the M shell are taken into account by the vacancy transfer coefficients n_{KL} and \bar{n}_{LM} , the expressions for the emission intensities are given by:

$$\begin{aligned}
 I_\gamma &= (1 - \alpha') P_g \\
 I_{ce} &= \alpha'_t P_g \\
 I_{X_K} &= \alpha'_K \omega_K P_g \\
 I_{A_K} &= \alpha'_K (1 - \omega_K) P_g \\
 I_{X_L} &= (\alpha'_K n_{KL} + \alpha'_L) \bar{\omega}_L P_g \\
 I_{A_L} &= (\alpha'_K n_{KL} + \alpha'_L) (1 - \bar{\omega}_L) P_g \\
 I_{X_M} &= (\alpha'_K n_{KL} \bar{n}_{LM} + \alpha'_L \bar{n}_{LM}) + \alpha'_M \bar{\omega}_L P_g \\
 I_{A_M} &= (\alpha'_K n_{KL} \bar{n}_{LM} + \alpha'_L \bar{n}_{LM}) + \alpha'_M (1 - \bar{\omega}_M) P_g
 \end{aligned}$$

In these equations ω_K , $\bar{\omega}_L$, $\bar{\omega}_M$, are the K-shell, the mean L-shell and the mean M-shell fluorescence yields and $\alpha'_i = \alpha_i / (1 + \alpha_t)$ are the reduced conversion coefficients ($i = K, L, M, t$). The equations containing $\bar{\omega}_L$, $\bar{\omega}_M$, are approximations. A more detailed description would require taking into account the processes occurring in the subshells. As the data necessary for this are usually known with less accuracy, there is often no advantage in comparison to the use of the above approximations. If there are several transitions, each contributes to the emission of X-ray quanta and Auger electrons and the contributions have to be summed up when the total emission intensities characterizing the disintegration of the studied nuclide are calculated.

Chapter 4

INTERACTION OF RADIATIONS WITH MATTER

This subject, for which the reader is advised to refer to basic texts [55Ev R, 68Si K], is raised only for two phenomena which occur in the vicinity of the radioactive source, and which have a direct effect on the intensity of the radiations detected.

4.1 External bremsstrahlung

The external bremsstrahlung radiation is derived from the slowing down of electrons in the Coulomb field of nuclei in the absorbing medium surrounding the β -emitting source. The average energy of this photon radiation is about $1,4 \times 10^{-4} Z E_{max}^2$, where E_{max} is the maximum energy of the β radiation, provided that it is totally absorbed in the material with atomic number Z (thick source).

The energy spectrum of this bremsstrahlung extends from 0 to E_{max} and decreases very roughly as $1/E$. Since the intensity of this radiation is a function of the thickness and nature of the material crossed, it is impossible to account for it in the tables. However, its influence on some γ and X-ray spectra can be assessed.

4.2 Annihilation-in-Flight of Positrons

A positron can annihilate with an electron of the traversed medium before a complete slowing down. The two emitted photons have energies different from $m_0 c^2$ as obtained from annihilation at rest [54Ge J]. In γ -ray spectrometry, this phenomenon has the effect of removing, from the 511 keV peak, a fraction of the annihilation photons, giving rise to a continuous photon spectrum. To take this effect into account the energy of this line is given as 511 keV without uncertainty as the energy is not exactly 511 keV. The magnitude of this effect depends on the material in which the β^+ are stopped, so the intensity of the 511 keV ray (I_{511}) does not have a unique value.

Several factors must be considered if one wants to use the measured intensity of the observed 511 keV peak to determine the activity of a radioactive source. First, all the β^+ particles need to be stopped near the source in an absorber of known material. Second, the point source efficiency at 511 keV must be corrected for the finite volume in which the β^+ annihilation occurs [80Sh K].

Chapter 5

DOSIMETRIC DATA

Administration of a radiopharmaceutical leads to the irradiation of various organs and tissues, which means that a prior estimate of the absorbed doses delivered to these parts of the body is necessary in order to evaluate the risks run by the individual concerned. These dose values are determined by physical and biological parameters and calculated from a computing model established by the “Medical Internal Radiation Dose Committee” [76Lo R], of the American Society of Nuclear Medicine. This model is based on the concepts of target region, in which the mean absorbed dose \bar{D} is estimated, as well as the source region from which the radiation arises.

The basic equation used for the calculations is [11ICRU]:

$$\bar{D}(r_k \leftarrow r_h) = \widetilde{A}_h \sum_i \Delta_i \frac{\varphi_i(r_k \leftarrow r_h)}{m_k} \quad (5.1)$$

where:

\widetilde{A}_h is the total number of nuclear transformations undergone by the radionuclide in the source region r_h (which denotes a volume) during the time interval considered. \widetilde{A}_h depends on the physical half-life of the radionuclide and on the activity delivered, as well as on biological parameters relative to the distribution in time of the radionuclide within the source region;

m_k is the mass of the target region r_k ;

Δ_i is the mean energy of i-type particles emitted by nuclear transformations of the radionuclide in question, obtained from the expression:

$$\Delta_i = n_i E_i \quad (5.2)$$

with:

n_i : the mean number of i-type particles emitted per nuclear transformation of the radionuclide.

E_i : the mean energy (other than the rest energy) of i-type particle.

$\varphi_i(r_k \leftarrow r_h)$, absorbed fraction, is the fraction of the energy E_i , of each particle emitted in the source region r_h , which is given, in mean, to the target region r_k .

A special case is that where the regions r_k and r_h are identical, exactly homogeneous, of constant density and in which the radionuclide is uniformly distributed. The general expression then reduces to the simplified form:

$$D_{eq} = \widetilde{C} \sum_i \Delta_i \quad (5.3)$$

where:

D_{eq} is the equilibrium dose,

$\tilde{C} = \tilde{A}/m$ the number of nuclear transformations per unit mass.

In this case the equilibrium results from the fact that at each point of the region the energy emitted is equal to that imparted to the medium.

The purely physical data considered in this Table and needed for calculations from the general expression (5.1) are:

- the radionuclide half-life, one of the parameters involved in \tilde{A}_h ,
- n_i , mean number of i-type particles per nuclear transformation,
- E_i , mean energy of each i-type particle,
- Δ_i , product of n_i and E_i .

The units of the various terms of equations (5.1) and (5.2) are given in the table below:

Term	SI Unit [81 SI]	Old unit and value in SI
\bar{D}	Gy	1 rad = 10^{-2} Gy
\tilde{A}_h	Bq.s = 1	1 μ Ci.h = $1,332 \times 10^8$ Bq.s
m_k	kg	1 g = 10^{-3} kg
n_i	Bq $^{-1}$.s $^{-1}$	1 g.rad. μ Ci $^{-1}$.h $^{-1}$.MeV $^{-1}$ = 1/2,134
E_i	J	1 MeV = $1,6022 \times 10^{-13}$ J
Δ_i	J.Bq $^{-1}$.s $^{-1}$	1 g.rad. μ Ci $^{-1}$.h $^{-1}$ \simeq $7,507 \times 10^{-14}$ J
φ_i	1	1

Chapter 6

RULES FOR EVALUATION AND COMPILATIONS

Two main groups of data sources are used to obtain the recommended data:

- specific values evaluated from all available original publications,
- compiled data already evaluated by specialists (for example: Q -values), if a new experimental value exists, it may be taken into account. In this case, the corresponding reference is mentioned in the reference list of this radionuclide.

6.1 Rules for evaluation

All intermediate stages in the compilation and evaluation of a decay parameter are not presented in detail in order to avoid unnecessary complexity. The main stages comprise the following:

- critical analysis of published results and, if necessary, correction of these results to account for more recent values hitherto unavailable to the original experimentalists; as a rule, results without associated uncertainties are discarded, and the rejection of values is documented;
- data obtained through private communications are only used when there is no published article available;
- adjustments may be made to the reported uncertainties;
- only one result, generally the latest one, is taken into account per laboratory (or author);
- recommended values are derived from an analysis of all available measurements (or theoretical considerations), along with the standard deviations corresponding to the 1σ confidence level.

6.1.1 Evaluation of uncertainties

Definitions by the “Guide to the expression of uncertainty in measurement” [93 GUM]:

Uncertainty (of measurement): a parameter, associated with the result of a measurement, that characterizes the dispersion of the values that could reasonably be attributed to the measurand.

Standard uncertainty: uncertainty of the result of a measurement expressed as a standard deviation.

Type A evaluation (of uncertainty): method of evaluation of uncertainty by the statistical analysis of a series of observations.

Type B evaluation (of uncertainty): method of evaluation of uncertainty by means other than the statistical analysis of a series of observations.

The uncertainties given by authors are re-evaluated by combining the standard uncertainties σ_A and σ_B by using the general law of variance propagation:

$$u_c = \sqrt{\sigma_A^2 + \sigma_B^2} \quad (6.1)$$

where: u_c : combined standard uncertainty
 σ_A : type A standard deviation
 σ_B : type B standard uncertainty

When the authors give insufficient information about their uncertainty calculations, the combined uncertainty u_c , may be estimated by the evaluator, based on his knowledge of the measurement method(s).

6.1.2 Determination of the best value and of the associated uncertainty

Case of results obtained by one author using one method

In this case the procedure is normally carried out by the author himself. Sometimes only the final result, for the mean value and the combined standard uncertainty, is given in the original publication. If details are known, the procedure is the following:

If there are n individual values a_i ($i = 1 \cdots n$), the best value is the arithmetical mean:

$$\bar{a} = \sum_{i=1}^n \frac{a_i}{n} \quad (6.2)$$

with type A standard deviation:

$$\sigma_A(\bar{a}) = \left[\frac{\sum_i (a_i - \bar{a})^2}{n(n-1)} \right]^{1/2} \quad (6.3)$$

If there are m contributions σ_{Bj} ($j = 1 \cdots k$) to the type B standard uncertainty that are independent of each other:

$$\sigma_B(\bar{a}) = \left[\sum_{j=1}^m \sigma_{Bj}^2 \right]^{1/2} \quad (6.4)$$

Combined standard uncertainty is:

$$u_c(\bar{a}) = \sqrt{\sigma_A^2(\bar{a}) + \sigma_B^2(\bar{a})} \quad (6.5)$$

Recommended value is:

$$a = \bar{a} \pm u_c(\bar{a}) \quad (6.6)$$

Case of results obtained by several authors employing the same method

For n individual values \bar{a}_i ($i = 1 \dots n$) having a standard deviation σ_{Ai} and a type B uncertainty σ_{Bi} the best value is obtained by taking the mean weighted by the inverse of the variances.

$$\bar{a} = \frac{\sum_j (\bar{a}_j / \sigma_{Aj}^2)}{\sum_j (1 / \sigma_{Aj}^2)} \quad (6.7)$$

The associated values u_c , σ_A , σ_B are:

$$\sigma_A(\bar{a}) = \left[\sum_i (1 / \sigma_{Ai}^2) \right]^{-1/2} \quad (6.8)$$

$\sigma_B(\bar{a}) = \sum_i (\sigma_{Bi})_{min}$ or $\sigma_B(\bar{a}) = \sqrt{\sum_i (\sigma_{Bi})_{min}^2}$ or $\sigma_B(\bar{a}) = (\sigma_B)_{min}$ depending on the individual case. However, $\sigma_B(\bar{a})$ cannot be less than the smallest σ_{Bi} .

Finally σ_A and σ_B are combined quadratically:

$$u_c(\bar{a}) = \sqrt{\sigma_A^2(\bar{a}) + \sigma_B^2(\bar{a})} \quad (6.9)$$

The recommended value is:

$$a = \bar{a} \pm u_c(\bar{a}) \quad (6.10)$$

Case of results obtained by different methods

When different measurement techniques have been applied, a weighted average is calculated using the combined uncertainties of the individual values as weights.

For n independent values a_i , each with a combined standard uncertainty u_{ci} ; a weight p_i proportional to the inverse of the square of the individual u_{ci} can be assigned to each value.

$$a_w = \frac{\sum_{i=1}^n p_i a_i}{\sum_{i=1}^n p_i} \quad (6.11)$$

where the weights are $p_i = 1/u_{ci}^2$.

An internal uncertainty can be assigned to the mean value [93Wi G, 85Zi P]:

$$\sigma_{int}(a_w) = \left[\sum_i (1/u_{ci}^2) \right]^{-1/2} \quad (6.12)$$

The internal variance $\sigma_{int}^2(a_w)$ is the expected uncertainty of the mean, based on the individual *a priori* variances $u_{c_i}^2$ (by uncertainty propagation).

The external uncertainty is also calculated:

$$\sigma_{ext}(a_w) = \left[\frac{\sum (a_i - a_w)^2 / u_{c_i}^2}{(n-1) \sum (1/u_{c_i}^2)} \right]^{1/2} \quad (6.13)$$

The external variance $\sigma_{ext}^2(a_w)$ includes the scatter of the data and is based on the amount by which each a_i deviates from the mean when measured as a fraction of its given uncertainty u_{c_i} .

A measure of the consistency of the data is given by the ratio [93Wi G, 85Zi P]:

$$\sigma_{ext}/\sigma_{int} = \sqrt{\chi^2/(n-1)} \quad (6.14)$$

If this ratio is significantly greater than unit, at least one of the input data most probably has an underestimated u_{c_i} which should be increased, or some data must be rejected.

Statistical rejection of data

The rejection of data is based on the Chauvenet's criterion [76Ch W, 72De D] that identifies statistical outliers.

In a set of n individual values a_i ($i = 1 \dots n$), having an arithmetical mean value of \bar{a} and a standard deviation of σ , a measured value a_k is identified as outlier if:

$$a_k > \bar{a} + \sigma \times C \quad (6.15)$$

where:

- C : depends on the number of measurements n and is calculated assuming that a_i ($i = 1 \dots n$) follows a Gaussian distribution.
- a_k : is rejected if the probability to obtain such value is less than 50 %.

Determination of the adopted value and of the associated uncertainty

After a possible rejection of data, a critical value of $\chi^2/(n-1)$ at 1 % confidence level is used as a practical test for discrepant data. The following table lists the critical values of $\chi^2/(n-1)$ (critical) for an increasing degree of freedom $\nu = N - 1$ [69Be P].

ν	$\chi^2/(n-1)$ (critical)	ν	$\chi^2/(n-1)$ (critical)
1	6,6	12	2,2
2	4,6	13	2,1
3	3,8	14	2,1
4	3,3	15	2,0
5	3,0	16	2,0
6	2,8	17	2,0
7	2,6	18 - 21	1,9

ν	$\chi^2/(n-1)$ (critical)	ν	$\chi^2/(n-1)$ (critical)
8	2,5	22 - 26	1,8
9	2,4	27 - 30	1,7
10	2,3	> 30	$1 + 2,33 \sqrt{2/\nu}$
11	2,2		

If $\chi^2/(n-1) \leq \text{critical } \chi^2/(n-1)$ the recommended value is given by:

$$a = a_w \pm \sigma_{int}(a_w) \quad (6.16)$$

If $1 < \chi^2/(n-1) > \text{critical } \chi^2/(n-1)$, the method of limitation of the relative statistical weight [85Zi P, TECDOC] is recommended when there are three or more values; uncertainty of a value contributing more than 50 % to the total weight is increased to reduce its contribution to 50 %. The weighted and unweighted average and $\text{critical } \chi^2/(n-1)$ are then recalculated:

If $\chi^2/(n-1) \leq \text{critical } \chi^2/(n-1)$ the recommended value is given by:

$$a = a_w \pm \max[\sigma_{int}(a_w), \sigma_{ext}(a_w)] \quad (6.17)$$

If $\chi^2/(n-1) > \text{critical } \chi^2/(n-1)$, the weighted or unweighted mean is chosen, depending on whether or not the uncertainties of the average values make them overlap with each other. If overlap occurs, the weighted average is recommended; otherwise the unweighted average is chosen. In either case, the uncertainty can be increased to cover the most accurate value.

Parameters evaluated according to these procedures and rules include half-lives, energies and number of emitted particles, and some experimental internal conversion coefficients. All remaining data given in these tables are taken from compilations or deduced from calculations.

6.1.3 Balanced decay schemes

All the probabilities for transitions correspond to balanced schemes and permit the formulation of a fully consistent set of values. This balance implies obvious relationships such as the following:

- The sum of the transition probabilities for all the decay transitions (α , β , γ , ϵ) cut by an horizontal plan at each level of a decay scheme is equal to 1 (or 100 %). This ruling is also valid for the highest energies of the decay scheme where only β (or α , ϵ) transitions are relevant, and for the lowest levels of the decay scheme, where sometimes only γ transitions are involved;
- for an excited nuclear level, the sum of the transition probabilities feeding the level (α , β , γ , ϵ) is equal to the sum of the transition probabilities depopulating this level;

- when the relative emission probabilities $P(rel)_{\gamma_i}$ of the γ emission intensities are known with respect to one of them, such as γ_k , and no feeding to the ground state from α , β , and ϵ transitions is assumed, then, all the absolute γ emission intensities $I(abs)_{\gamma_i}$ can be calculated from the equation:

$$I(abs)_{\gamma_i} = P(rel)_{\gamma_i} \times N \quad (6.18)$$

where N is the “normalization factor” and is determined by:

$$N \sum_i P(rel)_{\gamma_i} (1 + \alpha_{t_i}) = 1 \quad (6.19)$$

where the sum covers the γ -transitions feeding the ground state.

6.2 Compilations

6.2.1 β and electron capture transitions

Depending on the individual radionuclide, β -transition energies are either evaluated on the basis of experimental data (maximum β energies), or derived from the atomic mass differences obtained from the tabulations of AUDI, WAPSTRA, THIBAULT [03Au G] and the nuclear level energies. The average β energies are generally computed [ENSDF.], and the values of $\lg ft$ are calculated from the tables of GOVE and MARTIN [72Go N], as well as ϵ/β^+ when necessary.

Electron-capture transition energies are derived from the atomic mass differences and the nuclear level energies. Capture probabilities in the electronic shells, P_K , $P_L \dots$ are calculated from equations (3.37) to (3.49) where the ratios of the radial wave function components of the electron [69Su Y, 69Be H, 70Ma M], and the corrective terms for exchange $X^{L/K}$ [63Ba J, 70Fa J, 70Su Y, 70Va E, 98Sc E] are derived from tables.

6.2.2 γ transitions

- Internal conversion coefficients of pure transitions are evaluated and compared with theoretical values [02Ba I, 08Ki T], which are generally preferred, except for some specific cases, e.g. when there is a penetration effect.

The theoretical values are deduced from the tables of BAND *et al.* [02Ba I], they are based on the Dirac-Fock calculations using the so called “Frozen Orbital” approximation which takes into account the effect of atomic vacancies created in the conversion process [08Ki T].

The uncertainties on these theoretical values are deduced from a comparison between theoretical values and experimental ones [08Ki T, 02Ra S].

- Internal conversion coefficients of multipole transitions (e.g. M1 + E2) are derived from tables of theoretical data, according to:

$$\alpha_{T(M1+E2)} = (1 - \delta^2) \alpha_{T(M1)} + \delta^2 \alpha_{T(E2)} \quad (6.20)$$

Where the mixing ratio δ is evaluated from experimental values.

- α_π coefficients are interpolated from theoretical values [79Sc P, 96Ho C], the uncertainties are assumed to be 1,4 % [08Ki T].

6.2.3 Level spins and parities

Level spins and parities are usually extracted from the Nuclear Data Sheets [NuDaSh].

6.2.4 Constants of the atomic shell

The K-shell fluorescence yields ω_K and their uncertainties are taken from the evaluation of BAMBYNEK *et al.* [72Ba W, 84Ba W, 91Ba W] with uncertainties ranging from 1 % ($Z > 35$) to 10 % ($Z = 5$), and from subsequent experimental results.

Mean L-shell fluorescence yields $\bar{\omega}_L$ are taken from the evaluation of SCHÖNFELD *et al.* [95Sc E]. This evaluation includes both experimental [81Mi I, 87Co D, 89Hu J] and theoretical values [93Pu S]. The relative uncertainties are lower than 4 % (for $Z > 29$).

Mean M-shell fluorescence yields $\bar{\omega}_M$ are obtained from the fitting of experimental data by HUBBELL [89Hu J, 94Hu J].

Relative X-ray emission rates $K\beta/K\alpha$ are taken from SCHÖNFELD *et al.* [95Sc E], and the $K\alpha_1/K\alpha_2$ from the theoretical values of SCOFIELD [74Sc J]. Uncertainties are assumed to be of the order of 1 %.

X-ray radiation energies are taken from the tables of BEARDEN [67Be J] and DESLATTES [03De R].

Relative emission probabilities of K-Auger electron groups are deduced from the X-ray ratio [96Sc E] with uncertainties of the order of 3 %.

Energies of the K and L-Auger electrons are taken from the table of LARKINS [77La F].

The mean number of vacancies created in the L shell (from one K hole) n_{KL} and in the M shell (from one L hole) \bar{n}_{LM} are estimated from the previous values.

6.2.5 m_0c^2 energy

The m_0c^2 energy is taken as 510,998 928 (11) keV given by the CODATA Group (2010) [10Mo P].

Chapter 7

CONTENTS OF THE TABLES

Each section devoted to a radionuclide or to a radioactive series contains the following:

- decay scheme(s),
- half-lives, Q -values, atomic shell data,
- tables of probabilities of different transitions,
- tables of intensities of different emissions,
- major production methods,
- references employed in the evaluation.

7.1 Radioactive series

When two or more radionuclides compose a radioactive series, the nuclear and atomic data are presented separately for each. If the half-life of the daughter(s) is shorter than that of the parent, so that an equilibrium can be achieved, the emission intensities of the different radiations are given for all the radionuclides involved, the equilibrium being reached. Moreover, if the half-life of the daughter is sufficiently long to permit its use after separation, it is covered by a separate section (e.g. $^{140}\text{Ba} - ^{140}\text{La}$). This rule also applies if the daughter is an isomer (e.g. $^{125}\text{Sb} - ^{125\text{m}}\text{Te}$).

Furthermore, if a radionuclide possesses an isomer of specific interest, it is covered by a separate section (e.g. $^{58\text{m}}\text{Co}$, $^{60\text{m}}\text{Co}$).

7.2 Decay scheme

A decay scheme includes the following information:


- half-lives of ground states, together with those of excited levels longer than 10^{-12} s,
- disintegration energies,
- energy, spin and parity of levels,
- transitions listed by the index: initial level, final level. The levels are arbitrarily designated by a number from 0 for the ground state to n for the n^{th} excited level.

The transition energies are not shown on the decay schemes. The α , β and electron capture and γ transition probabilities are given for 100 disintegrations. The gamma emission intensities are given following 100 disintegrations.


Intervals between levels are roughly proportional to energy differences.

Transitions with probabilities over 0,5 are shown by heavy lines, as used for the ground states. Those between 0,05 and 0,5 in medium-thick lines. Those below 0,05 are shown by fine lines.


The decay scheme diagram is, as a rule, similar to that of other tables [NuDaSh].




β decay.




Electron capture decay.




β^+ decay.



α decay.



Q values of disintegration energies, Q^- being shown on the right of the schemes, Q^+ and Q_α on the left.



The upper level has an energy of 150 keV and a half-life of $1,4 \times 10^{-9}$ s. It de-excites by the emission of a γ photon or an internal conversion electron.

<u>2⁺</u>	<i>Spin and parity are considered as certain.</i>
<u>(4⁺)</u>	<i>Spin and parity are uncertain.</i>
<u>(4⁺)</u>	<i>Spin is uncertain, parity is considered as certain.</i>
<u>2⁽⁺⁾</u>	<i>Parity is uncertain, spin is considered as certain.</i>
<u>(3/2, 1/2)⁻</u>	<i>Parity is considered as certain, and one of the spins as certain.</i>

7.3 Tables

7.3.1 Notation

Following the Guide to the expression of uncertainty in measurement [93 GUM], the values and their uncertainties are reported as:

$$E = 100,021\,47(35) \text{ keV}$$

where the number in parentheses, which is the numerical value of the combined standard uncertainty u_c ($k = 1$), refers to the corresponding last digits of the quoted result.

7.3.2 Nuclear data

The nuclear data comprise the following:

- the half-life of the radionuclide and possibly those of the daughter radionuclides,
- disintegration energies,
- α transitions, β transitions and electron capture transitions are given in order of increasing energy. The first column designates the transition. The remaining columns indicate the energy, the transition probability, the α transition hindrance factor, the nature, the $\lg ft$ for β transitions and, for electron capture transitions, the probabilities P_{ϵ_K} , P_{ϵ_L} , P_{ϵ_M} of capture in the atomic shells.

- γ transitions are also given by order of increasing energy. The first column designates the transition. The remaining columns indicate the energy, probability, multipolarity, internal conversion coefficients α_K , α_L , the total conversion coefficient α_t and the coefficient of creation of the internal pair α_π if any.

Remarks:

A multipolarity such as M1 (+ E2) indicates that the percentage of E2 is very low or is uncertain.

A multipolarity such as M1 or E2 indicates a pure γ transition of one type.

7.3.3 Atomic data

Atomic data include the following:

- fluorescence yields ω_K , $\bar{\omega}_L$, $\bar{\omega}_M$, n_{KL} , \bar{n}_{LM} ,
- energies and relative probabilities of X_K lines,
- energies of X_L lines. For radionuclides with $Z \leq 52$, only the minimum and maximum energies are indicated,
- energies and relative emission probabilities of K-Auger electrons,
- minimum and maximum energies of L-Auger electrons. The minimum energy is that of the $L_3 M_1 M_1$ etc. transition. The maximum energy is that of the $L_1 MM$ transition for $Z \leq 32$, of the $L_1 NN$ transition for $33 \leq Z \leq 37$, of the $L_1 NO$ transition for $38 \leq Z \leq 49$ and of the $L_1 OO$ transition for $Z \geq 50$.

7.3.4 Intensities of radiations

Emission intensities are always related to 100 nuclear transformations of the radionuclide and are given in order of increasing energy.

Isomeric transitions

When a radionuclide with half-life T_1 has an isomer daughter with half-life T_2 lower than T_1 , the emission intensities I_{γ_i} and I_{ce_i} are related to the probability P_{g_i} of the isomeric transition by:

$$I_{\gamma_i} + I_{ce_i} = \frac{T_1}{T_1 - T_2} P_{g_i} \left[1 - e^{-\ln 2 \frac{T_1 - T_2}{T_1 T_2} t} \right] \quad (7.1)$$

For $t \geq 10 T_2$, the parent and daughter are practically in equilibrium:

$$I_{\gamma_i} + I_{ce_i} = \frac{T_1}{T_1 - T_2} P_{g_i} \quad (7.2)$$

Note that the isomeric transition probability P_{g_i} is the probability that the disintegration of the parent radionuclide is followed by this transition. It is independent of the time and cannot exceed 1 (or 100 %).

7.4 Main production modes

The production processes quoted are those which are most widely employed. The impurities mentioned are those generally encountered, with the exception of accidental contaminations. The cross section, given only for thermal neutron activation, is taken from the tables of MUGHABGHAB *et al.* [81Mu S, 06Mu S]. The useful data for activation with charged particles (cross section as a function of particle energy) may be derived from tables or publications [89He R].

7.5 References used

The references quoted are those which were used in the evaluations, with the extracted quantities in square brackets.

7.6 Spectra

Spectra are not given in the new computerized version of these tables, they can be found in the Gamma-Ray Spectrum Catalog [98Cat.].

7.7 Publications

Various volumes of the “Table of Radionuclides” were published [87 TR1, 99 TR2]. A new series of volumes have been published under the auspices of the BIPM as Monographie-5 [04TRM1, 06TRM2, 08TRM3, 10TRM4, 11TRM5]. Each evaluation includes a comment file published in a separate issue.

New publications or updates are made available in the following web site:

<http://www.nucleide.org/NucData.htm>

A CD Rom, called “Nucleide” has also been published [CD NUC], it includes all Tables, Comments and various facilities.

For spectrometry purposes, a web application has been developed. It presents all α , X and γ emissions of a given nuclide and provides multi-criteria searching tools. It can be found at the following web address:

<http://laraweb.free.fr>

A pocket table containing the main characteristics of a decay is also provided [07Mini].

REFERENCES

- [40Mo P] P. MORRISSON, L.I. SCHIFF, Phys. Rev. 58 (1940) 24
- [49Ch C] C.S. CHANG, D.L. FALKOFF, Phys. Rev. 76 (1949) 365
- [51We V] V.F. WEISSKOPF, Phys. Rev. 83 (1951) 1073
- [53Bi L] L.C. BIEDENHARN, M.E. ROSE, Rev. Mod. Phys. 25 (1953) 729
- [53Pr H] H. PRIMAKOFF, F.T. PORTER, Phys. Rev. 89 (1953) 930
- [54Ge J] J.B. GERHART, B.C. CARLSON, R. SHERR, Phys. Rev. 94, 4 (1954) 917
- [55Ev R] R.D. EVANS, The Atomic Nucleus (Mc Graw-Hill, New York) (1955)
- [60Bo R] R. BOUCHEZ, P. DEPOMMIER, Repts. Progr. in Phys. 23 (1960) 395
- [60Mi H] H.V. MICHEL, Report U.C.R.L. 9229 (1960)
- [63Ba J] J.N. BAHCALL, Phys. Rev. 129 (1963) 2683, Phys. Rev. 132 (1963) 362 and Nucl. Phys. 71 (1965) 267
- [64Hy E] E.K. HYDE, I. PERLMAN, G.T. SEABORG, The Nuclear Properties of the Heavy Elements (ed. Prentice Hall inc.) (1964) Vol. 1
- [65Fi R] R.W. FINK, R.C. JOPSON, H. MARK, C.D. SWIFT, Rev. Mod. Phys. 38 (1965) 513
- [66Go N] N.B. GOVE, Nuclear Spin Parity Assignment (ed. N.B. Gove Academic Press N.Y.) (1966) p. 83
- [66Ko E] E.J. KONOPINSKY, The Theory of Beta Radioactivity, (Oxford Univ. Press) (1966)
- [66Sc H] H.F. SCHOPPER, Weak Interactions and Nuclear Beta Decay, (North-Holland Publ. Co. Amsterdam) (1966)
- [67Be J] J.A. BEARDEN, Rev. Mod. Phys. 39 (1967) 78 and 125
- [68Le H] H.J. LEISI, Proceedings of the Conference on the Electron Capture and Higher Order Process in Nuclear Decays (ed. D. Berenyi), Vol. 3 (1968) p. 368
- [68Pe B] B.I. PERSSON, Proceedings of the Conference on the Electron Capture and Higher Order Processes in Nuclear Decays (ed. D. Berenyi) Vol. 2 (1968) 142

- [68Ra J] J.O. RASMUSSEN, in 'Alpha, Beta and Gamma Ray spectroscopy' (ed. K. Siegbahn, North Holland Publishing Company) (1968)
- [68Si K] Alpha, Beta and Gamma Ray Spectroscopy (ed. K. Siegbahn)(North-Holland Publishing Company) (1968) Vol. 1 & Vol. 2
- [68Wi J] J.C. WIDMAN, J. MANTEL, N.M. HORWITZ, E. R. POWSNER, Int. J. Apl. Radiat. Isot. 19 (1968) 1
- [68Wi R] R. WILSON in Alpha, Beta and Gamma Ray Spectroscopy (ed. K. Siegbahn)(North-Holland Publishing Company) (1968) Vol. 2, p. 1557
- [68Zy J] J. ZYLICZ, Proceedings of the Conference on the Electron Capture and Higher Order Processes in Nuclear Decays (ed. D. Berenyi) Vol. 2 (1968) 123
- [69Be H] H. BEHRENS, J. JANECKE, in Landolt-Bornstein: Numerical Data and Functional Relationships in Science and Technology, New Series Group 1, Vol. 4 - Numerical Tables for Beta Decay and Electron Capture (Springer Verlag Berlin)(1969).
- [69Be P] P.R. BEVINGTON, Data Reduction and Error Analysis for the Physical Sciences, McGraw-Hill, Appendix C-4 (1969)
- [69St P] P. STEPHAS, Phys. Rev. 186 (1969) 1013
- [69Su Y] Y.P. SUSLOV, Bull. Acad. Sci. USSR Fiz. 33 (1969) 74
- [70Fa J] J.A. FAESSLER, E. HUSTER, O. KRAFFT, F. KRAHN, Z. Phys. 238 (1970) 352
- [70Ma M] M.J. MARTIN, P.H. BLICHERT-TOFT, Nuclear Data Tables A8 (1970) 1
- [70Su Y] Y.P. SUSLOV, Bull. Acad. Sci. USSR, Ser. Fi. 34 (1970) 91
- [70Va E] E. VATAI, Nucl. Phys. A156 (1970) 541
- [71Ch M] M.H. CHEN, B. CRASEMANN, V.O. KOSTROUN, Phys. Rev. A4 (1971) 1
- [71Cr B] B. CRASEMANN, M.H. CHEN, V.O. KOSTROUN, Phys. Rev. A4 (1971) 2161
- [71Ma E] E.J. MAC GUIRE, Phys. Rev. A3 (1971) 587
- [72Ba W] W. BAMBYNEK, *et al.*, Rev. Mod. Phys. 44 (1972) 716
- [72De D] D. DE SOETE, *et al.*, Neutron Activation Analysis, Wiley (1972) 505
- [72Em G] G.T. EMERY, Ann. Rev. Nucl. Sci. 22 (1972) 165
- [72Go N] N.B. GOVE, M.J. MARTIN, Nuclear Data Tables A10 (1971) 205
- [72Sh S] Proceedings of the International Conference on Inner Shell Ionization Phenomena and Future Applications CONF-720404, Atlanta (1972) Vol. 3. S. SHIMIZU p.2050 and A.J. MORD p.2079 and J. LAW, J.L. CAMPBELL p. 2110
- [73Cr B] B. CRASEMANN, Nucl. Instr. and Meth. 112 (1973) 33
- [73Ra S] S. RAMAN, N.B. GOVE, Phys. Rev. C7 (1973) 1995

- [73Sp A] A. SPERNOL, E. DE ROOST, M. MUTTERER, Nucl. Instr. and Meth. 112 (1973) 169
- [74Sc J] J.H. SCOFIELD, Atomic Data and Nuclear Data Tables 14 (1974) 121 and Phys. Rev. A9 (1974) 1041
- [74Sa S] S.I. SALEM, S.L. PANOSSIAN, R.A. KRAUSE, Atomic Data and Nuclear Data Tables 14 (1974) 91
- [75Pa H] H.C. PAULI, U. RAFF. A computer program for internal conversion coefficients and particles parameters. Computer Physics Communications 9 (1975) 392
- [76Be H] H. BEHRENS, L. SZYBISZ, Shapes of β spectra, ZAED, Physics Data 6-1 (1976)
- [76Ch W] W. CHAUVENET, 1876. Criterion for the rejection of doubtful observations. A manual of spherical and practical astronomy. Vol. 2, p 558, V.B. Lippincott & Co, Philadelphia
- [76Lo R] R. LOEVINGER, M. BERMAN, Mird-Pamphlet 1, revised (1976)
- [77La F] F.P. LARKINS, Atomic Data and Nuclear Data Tables 20 (1977) 313
- [78Po F] F.T. PORTER, M.S. FREEDMAN, J. Phys. Chem. Ref. Data Vol. 7,4 (1978) 1267
- [79Ch M] M.H. CHEN, B. CRASEMANN, H. MARK, Atomic Data and Nuclear Data Tables 24 (1979) 13
- [79Sc P] P. SCHLÜTER, G. SOFF, Atomic Data and Nuclear Data Tables 24 (1979) 509
- [80Ch M] M.H. CHEN, B. CRASEMANN, H. MARK, Phys. Rev. A21 (1980) 436
- [80Co N] N. COURSOL, Rapport CEA-R-5052 (1980)
- [80Sh K] K. SHIZUMA, S. HAMANAKA, Y. YOSHIKAWA, J. Phys. Soc. Japan, 49, 3 (1980) 847
- [81Mi I] I.V. MITCHELL, K.M. BARFOOT, Particle induced X-Ray Emission Analysis, Application to Analytical Problems, Nucl. Sci. Appl. (Harwood, UK) 1 (1981) 99-162
- [81Mu S] S.F. MUGHABGHAB, M. DIVADEENAM, N.E. HOLDEN, Neutron Cross Sections - Vol. 1, Academic Press (1981)
- [81 SI] Le Système International d'Unités, Bureau International des Poids et Mesures, Pavillon de Breteuil, Sèvres, France, 4e Edition (1981)
- [84Ba W] W. BAMBYNEK, Proc. Conf. on X-Ray and Inner Shell Processes in Atoms, Molecules and Solids, Aug. 20-24, Leipzig, Germany (1984)
- [85Zi P] P.L. ZIJP, Netherland Energy Research Foundation ECN, Petten, The Netherlands, Report ECN-179 (1985)
- [87Co D] D.D. COHEN, Nucl. Instr. Methods B22 (1987) 55

- [89He R] R. HEATON, H. LEE, P. SKENSVED, B.C. ROBERTSON. Nucl. Instr. Methods A276 (1989) 529.
- [89Hu J] J.H. HUBBELL, Bibliography and Current Status of K, L, and higher Shell Fluorescence Yields for Computation of Photon Energy Absorption Coefficients. NIST Internal Report 89-4144 (1989)
- [90Ji W] W. JITSCHIN, Progress in Measurements of L-Subshell Fluorescence, Coster-Kronig and Auger Yields, AIP Conf. Proc. (USA) 215 (1990) 408, 15. Int. Conf. on X-Ray and Inner Shell Processes (X-90), Knoxville TN, USA, 9-13 July 1990
- [91Ba W] W. BAMBYNEK, Reorganization of atomic shells after radioactive decay in Proc. of the Second Int. Summer School, Low-Level Measurements of man-made Radionuclides in the environment, La Rabida, Huelva, Spain, June 1990, World Scientific, Singapore (1991) 156-174
- [91Ca J] J.L. CAMPBELL, J.A. MAXWELL, W.J. TEESDALE, Phys. Rev. C 43,4 (1991) 1656
- [92 ISO] ISO 31, Quantities and Units, ISO Standards Handbook, CP 56, CH-1211 Genève 20, ISBN 92-67-10185-4 (1992)
- [93aPuS] S. PURI, D. MEHTA, B. CHAND, N. SINGH, P.N. TREHAN, Nucl. Instr. Methods B74 (1993) 347
- [93 GUM] Guide to the expression of uncertainty in measurement (1993) ISBN 92-67-20185-X, ISO Geneva.
- [93Pu S] S. PURI, D. MEHTA, B. CHAND, N. SINGH, P. N. TREHAN, X-Ray Spectrometry 22 (1993) 358
- [93Wi G] G. WINKLER, Data fitting and evaluation techniques for radioactive decay data in Proc. Symp. on Nuclear Data Evaluation Methodology, Oct 1992, Brookhaven Nat. Lab., U.S.A., World Scientific (1993)
- [94Hu J] J.H. HUBBELL, P.N. TREHAN, N. SINGH, B. CHAND, D. MEHTA, M.L. GARG, R.R. GARG, S. SINGH, S. PURI, J. Phys. Chem. Ref. Data Vol. 23,2 (1994) 339
- [95Sc E] E. SCHÖNFELD, H. JANSSEN, PTB-Report Ra-37 (1995)
- [96Ho C] C.R. HOFMANN, G. SOFF, Atomic Data and Nuclear Data Tables 63 (1996) 189
- [96Sc E] E. SCHÖNFELD, H. JANSSEN, Nucl. Instr. Meth A369 (1996) 527
- [98Cat.] GAMMA-RAY SPECTRUM CATALOG, CD-Rom (1998), INEEL, P.O.Box 1625, Idaho Falls, ID 83415, USA (<http://id.inel.gov/gamma/data1.html>)
- [98Sc E] E. SCHÖNFELD, Appl. Radiations Isot. 49,9 (1998) 1353
- [98Si B] B. SINGH, *et al.*, Nuclear Data Sheets 84,3 (1998) 487
- [00Mo P] P. J. MOHR, B. N. TAYLOR, Rev. Mod. Phys. 72,2 (2000) 351

- [02Ba I] I.M. BAND, M.B. TRZHASKOVSKAYA, M.A. LISTENGARTEN, Atomic Data and Nuclear Data Tables 81(2002)1
- [02Ra S] S. RAMAN, *et al.*, Phys. Rev. C66 (2002) 044312
- [03Au G] G. AUDI, A.H. WAPSTRA, C. THIBAULT. The 2003 Atomic Mass Evaluation, Nucl. Phys. A729 (2003) 337
- [03De R] R.D. DESLATTES, *et al.*, Rev. Mod. Phys. 75 (2003) 35
- [06Mu S] S.F. MUGHABGHAB, Atlas of Neutron Resonances, Elsevier Ed. (2006)
- [08Ki T] T. KIBÉDI, *et al.*, Nucl. Instr. and Meth. A 589 (2008) 202
- [10Mo P] P.J. MOHR, B.N. TAYLOR, D.B. NEWELL, The 2010 CODATA Recommended Values of the Fundamental Physical Constants, <http://physics.nist.gov/constants>, Web version 6.0 (2011)
- [11ICRU] I.C.R.U. Report 85, International Commission on Radiation Units and Measurements (2011)
- [11Mo X] X. MOUGEOT, M.M. BÉ, V. CHISTÉ, C. DULIEU, V. GOROZHANKIN, M. LOIDL, Radiocarbon (2011), LSC2010, Advances in Liquid Scintillation Spectrometry, 6-10 Sept. 2010, p. 249, ISBN 978-0-9638314-7-7
- [87 TR1] Table de Radionucléides. F. Lagoutine, N. Coursol, J. Legrand, ISBN-2-7272-0078-1. LMRI, 1982-1987, BP 52, 91 191 Gif-sur-Yvette Cedex, France.
- [99 TR2] Table de Radionucléides, Table of Radionuclides, Vol. 5. 1999, ISBN-2-7272-0200-8 and Comments ISBN-2-7272-0211-3 (1999) BNM-CEA/LPRI, BP 52, 91 191 Gif-sur-Yvette Cedex, France.
- [04TRM1] M.M. BÉ, E. BROWNE, V. CHECHEV, V. CHISTÉ, R. DERSCH, C. DULIEU, R.G. HELMER, D. McMAHON, A.L. NICHOLS, E. SCHÖNFELD, Table de Radionucléides, Monographie BIPM-5, Vol. 1 & 2 (2004), ISBN 92-822-2207-7 (set) and ISBN 92-822-2205-5 (CD), CEA/BNM-LNHB, 91191 Gif-sur-Yvette, France and BIPM, Pavillon de Breteuil, 92312 Sèvres, France.
- [06TRM2] M.M. BÉ, E. BROWNE, V. CHECHEV, V. CHISTÉ, R. DERSCH, C. DULIEU, R.G. HELMER, F.G. KONDEV, D. McMAHON, A.L. NICHOLS, E. SCHÖNFELD, Table of Radionuclides, Monographie BIPM-5, Vol. 3 (2006), ISSN 92-822-2204-7 (set), ISBN 92-822-2218-7 (Vol. 3) and ISBN 92-822-2219-5 (CD), CEA/LNE-LNHB, 91191 Gif-sur-Yvette, France and BIPM, Pavillon de Breteuil, 92312 Sèvres, France.
- [07Mini] Mini Table de Radionucléides (2007), CEA/LIST-LNHB, 91191 Gif-sur-Yvette, France, EDP Sciences Editeur, ISBN 978-2-86883-973-2
- [08TRM3] M.M. BÉ, E. BROWNE, V. CHECHEV, V. CHISTÉ, C. DULIEU, M. GALÀN, F.G. KONDEV, N. KUZMENKO, A. LUCA, A. PEARCE, X. HUANG, Table of Radionuclides, Monographie BIPM-5, Vol. 4 (2008), ISBN 92-822-2230-6 and ISBN 92-822-2231-4 (CD), CEA/LNE-LNHB, 91191 Gif-sur-Yvette, France and BIPM, Pavillon de Breteuil, 92312 Sèvres, France.

- [10TRM4] M.M. BÉ, E. BROWNE, V. CHECHEV, V. CHISTÉ, C. DULIEU, M. GALÀN, F.G. KONDEV, N. KUZMENKO, A. LUCA, A. ARINC, X. HUANG, A.L. NICHOLS, Table of Radionuclides, Monographie BIPM-5, Vol. 5 (2010), ISBN 978-92-822-2234-8 (Vol. 5) and 978-92-822-2235-5 (CD-Rom), CEA/LNE-LNHB, 91191 Gif-sur-Yvette, France and BIPM, Pavillon de Breteuil, 92312 Sèvres, France.
- [11TRM5] M.M. BÉ, V. CHISTÉ, C. DULIEU, X. MOUGEOT, V. CHECHEV, N. KUZMENKO, F.G. KONDEV, A. LUCA, M. GALÀN, A.L. NICHOLS, A. ARINC, A. PEARCE, X. HUANG, B. WANG, Table of Radionuclides, Monographie BIPM-5, Vol. 6 (2011), ISBN 978-92-822-2242-3 (Vol. 6) and 978-92-822-2243-0 (CD-Rom), CEA/LNE-LNHB, 91191 Gif-sur-Yvette, France and BIPM, Pavillon de Breteuil, 92312 Sèvres, France.
- [TECDOC] X-ray and gamma-ray standards for detector calibration, IAEA-TECDOC-619, Vienna (1991)
- [ENSDF.] Logft program, ENSDF Analysis and checking codes, Brookhaven National Laboratory
- [NuDaSh] NUCLEAR DATA SHEETS, National Nuclear Data Center, Brookhaven National Laboratory, Oak Ridge, Tennessee 37831-6371, USA.
- [CD NUC] M.M. BÉ, E. BROWNE, V. CHECHEV, V. CHISTÉ, C. DULIEU, M. GALÀN, R.G. HELMER, A. LUCA, X. MOUGEOT, N. KUZMENKO, A.L. NICHOLS, A. PEARCE, E. SCHÖNFELD, NUCLÉIDE, Table de Radionucléides sur CD-Rom, Version 3-2010, CEA/LIST-LNHB, 91191 Gif-sur-Yvette, France, EDP Sciences Editeur, ISBN 978-2-7598-0077-3